

Modern Physics

PART 6

At the end of the nineteenth century, many scientists believed that they had learned most of what there was to know about physics. Newton's laws of motion and his theory of universal gravitation, Maxwell's theoretical work in unifying electricity and magnetism, the laws of thermodynamics and kinetic theory, and the principles of optics were highly successful in explaining a variety of phenomena.

As the nineteenth century turned to the twentieth, however, a major revolution shook the world of physics. In 1900 Planck provided the basic ideas that led to the formulation of the quantum theory, and in 1905 Einstein formulated his brilliant special theory of relativity. The excitement of the times is captured in Einstein's own words: "It was a marvelous time to be alive." Both ideas were to have a profound effect on our understanding of nature. Within a few decades, these two theories inspired new developments and theories in the fields of atomic physics, nuclear physics, and condensed-matter physics.

In Chapter 39 we introduce the special theory of relativity. The theory provides us with a new and deeper view of physical laws. Although the concepts underlying this theory often violate our common sense, the theory correctly predicts the results of experiments involving speeds near the speed of light. In the extended version of this textbook, *Physics for Scientists and Engineers with Modern Physics*, we cover the basic concepts of quantum mechanics and their application to atomic and molecular physics, and we introduce solid-state physics, nuclear physics, particle physics, and cosmology.

You should keep in mind that, although the physics that was developed during the twentieth century has led to a multitude of important technological achievements, the story is still incomplete. Discoveries will continue to evolve during our lifetimes, and many of these discoveries will deepen or refine our understanding of nature and the world around us. It is still a "marvelous time to be alive." ■

◀ A portion of the accelerator tunnel at Fermilab, near Chicago, Illinois. The tunnel is circular and 1.9 km in diameter. Using electric and magnetic fields, protons and antiprotons are accelerated to speeds close to that of light and then allowed to collide head-on, in order to investigate the production of new particles. (Fermilab Photo)



Chapter 39

Relativity

CHAPTER OUTLINE

- 39.1 The Principle of Galilean Relativity
- 39.2 The Michelson–Morley Experiment
- 39.3 Einstein's Principle of Relativity
- 39.4 Consequences of the Special Theory of Relativity
- 39.5 The Lorentz Transformation Equations
- 39.6 The Lorentz Velocity Transformation Equations
- 39.7 Relativistic Linear Momentum and the Relativistic Form of Newton's Laws
- 39.8 Relativistic Energy
- 39.9 Mass and Energy
- 39.10 The General Theory of Relativity



▲ Standing on the shoulders of a giant. *David Serway, son of one of the authors, watches over his children, Nathan and Kaitlyn, as they frolic in the arms of Albert Einstein at the Einstein memorial in Washington, D.C. It is well known that Einstein, the principal architect of relativity, was very fond of children. (Emily Serway)*



Our everyday experiences and observations have to do with objects that move at speeds much less than the speed of light. Newtonian mechanics was formulated by observing and describing the motion of such objects, and this formalism is very successful in describing a wide range of phenomena that occur at low speeds. However, it fails to describe properly the motion of objects whose speeds approach that of light.

Experimentally, the predictions of Newtonian theory can be tested at high speeds by accelerating electrons or other charged particles through a large electric potential difference. For example, it is possible to accelerate an electron to a speed of $0.99c$ (where c is the speed of light) by using a potential difference of several million volts. According to Newtonian mechanics, if the potential difference is increased by a factor of 4, the electron's kinetic energy is four times greater and its speed should double to $1.98c$. However, experiments show that the speed of the electron—as well as the speed of any other object in the Universe—always remains less than the speed of light, regardless of the size of the accelerating voltage. Because it places no upper limit on speed, Newtonian mechanics is contrary to modern experimental results and is clearly a limited theory.

In 1905, at the age of only 26, Einstein published his special theory of relativity. Regarding the theory, Einstein wrote:

The relativity theory arose from necessity, from serious and deep contradictions in the old theory from which there seemed no escape. The strength of the new theory lies in the consistency and simplicity with which it solves all these difficulties¹

Although Einstein made many other important contributions to science, the special theory of relativity alone represents one of the greatest intellectual achievements of all time. With this theory, experimental observations can be correctly predicted over the range of speeds from $v = 0$ to speeds approaching the speed of light. At low speeds, Einstein's theory reduces to Newtonian mechanics as a limiting situation. It is important to recognize that Einstein was working on electromagnetism when he developed the special theory of relativity. He was convinced that Maxwell's equations were correct, and in order to reconcile them with one of his postulates, he was forced into the revolutionary notion of assuming that space and time are not absolute.

This chapter gives an introduction to the special theory of relativity, with emphasis on some of its consequences. The special theory covers phenomena such as the slowing down of moving clocks and the contraction of moving lengths. We also discuss the relativistic forms of momentum and energy.

In addition to its well-known and essential role in theoretical physics, the special theory of relativity has practical applications, including the design of nuclear power plants and modern global positioning system (GPS) units. These devices do not work if designed in accordance with nonrelativistic principles.

¹ A. Einstein and L. Infeld, *The Evolution of Physics*, New York, Simon and Schuster, 1961.

39.1 The Principle of Galilean Relativity

To describe a physical event, we must establish a frame of reference. You should recall from Chapter 5 that an inertial frame of reference is one in which an object is observed to have no acceleration when no forces act on it. Furthermore, any system moving with constant velocity with respect to an inertial frame must also be in an inertial frame.

There is no absolute inertial reference frame. This means that the results of an experiment performed in a vehicle moving with uniform velocity will be identical to the results of the same experiment performed in a stationary vehicle. The formal statement of this result is called the **principle of Galilean relativity**:

Principle of Galilean relativity

The laws of mechanics must be the same in all inertial frames of reference.

Let us consider an observation that illustrates the equivalence of the laws of mechanics in different inertial frames. A pickup truck moves with a constant velocity, as shown in Figure 39.1a. If a passenger in the truck throws a ball straight up, and if air effects are neglected, the passenger observes that the ball moves in a vertical path. The motion of the ball appears to be precisely the same as if the ball were thrown by a person at rest on the Earth. The law of universal gravitation and the equations of motion under constant acceleration are obeyed whether the truck is at rest or in uniform motion.

Both observers agree on the laws of physics—they each throw a ball straight up and it rises and falls back into their hand. What about the *path* of the ball thrown by the observer in the truck? Do the observers agree on the path? The observer on the ground sees the path of the ball as a parabola, as illustrated in Figure 39.1b, while, as mentioned earlier, the observer in the truck sees the ball move in a vertical path. Furthermore, according to the observer on the ground, the ball has a horizontal component of velocity equal to the velocity of the truck. Although the two observers disagree on certain aspects of the situation, they agree on the validity of Newton's laws and on such classical principles as conservation of energy and conservation of linear momentum. This agreement implies that no mechanical experiment can detect any difference between the two inertial frames. The only thing that can be detected is the relative motion of one frame with respect to the other.

Quick Quiz 39.1 Which observer in Figure 39.1 sees the ball's *correct* path? (a) the observer in the truck (b) the observer on the ground (c) both observers.




Figure 39.1 (a) The observer in the truck sees the ball move in a vertical path when thrown upward. (b) The Earth observer sees the path of the ball as a parabola.

Suppose that some physical phenomenon, which we call an *event*, occurs and is observed by an observer at rest in an inertial reference frame. The event's location and time of occurrence can be specified by the four coordinates (x, y, z, t) . We would like to be able to transform these coordinates from those of an observer in one inertial frame to those of another observer in a frame moving with uniform relative velocity compared to the first frame. When we say an observer is "in a frame," we mean that the observer is at rest with respect to the origin of that frame.

Consider two inertial frames S and S' (Fig. 39.2). The frame S' moves with a constant velocity \mathbf{v} along the common x and x' axes, where \mathbf{v} is measured relative to S. We assume that the origins of S and S' coincide at $t = 0$ and that an event occurs at point P in space at some instant of time. An observer in S describes the event with space–time coordinates (x, y, z, t) , whereas an observer in S' uses the coordinates (x', y', z', t') to describe the same event. As we see from the geometry in Figure 39.2, the relationships among these various coordinates can be written

$$x' = x - vt \quad y' = y \quad z' = z \quad t' = t \quad (39.1)$$

These equations are the **Galilean space–time transformation equations**. Note that time is assumed to be the same in both inertial frames. That is, within the framework of classical mechanics, all clocks run at the same rate, regardless of their velocity, so that the time at which an event occurs for an observer in S is the same as the time for the same event in S'. Consequently, the time interval between two successive events should be the same for both observers. Although this assumption may seem obvious, it turns out to be incorrect in situations where v is comparable to the speed of light.

Now suppose that a particle moves through a displacement of magnitude dx along the x axis in a time interval dt as measured by an observer in S. It follows from Equations 39.1 that the corresponding displacement dx' measured by an observer in S' is $dx' = dx - v dt$, where frame S' is moving with speed v in the x direction relative to frame S. Because $dt = dt'$, we find that

$$\frac{dx'}{dt'} = \frac{dx}{dt} - v$$

or

$$u'_x = u_x - v \quad (39.2)$$

where u_x and u'_x are the x components of the velocity of the particle measured by observers in S and S', respectively. (We use the symbol \mathbf{u} for particle velocity rather than \mathbf{v} , which is used for the relative velocity of two reference frames.) This is the **Galilean velocity transformation equation**. It is consistent with our intuitive notion of time and space as well as with our discussions in Section 4.6. As we shall soon see, however, it leads to serious contradictions when applied to electromagnetic waves.

Quick Quiz 39.2 A baseball pitcher with a 90-mi/h fastball throws a ball while standing on a railroad flatcar moving at 110 mi/h. The ball is thrown in the same direction as that of the velocity of the train. Applying the Galilean velocity transformation equation, the speed of the ball relative to the Earth is (a) 90 mi/h (b) 110 mi/h (c) 20 mi/h (d) 200 mi/h (e) impossible to determine.

The Speed of Light

It is quite natural to ask whether the principle of Galilean relativity also applies to electricity, magnetism, and optics. Experiments indicate that the answer is no. Recall from Chapter 34 that Maxwell showed that the speed of light in free space is $c = 3.00 \times 10^8$ m/s. Physicists of the late 1800s thought that light waves moved through a medium called the *ether* and that the speed of light was c only in a special, absolute frame




Figure 39.2 An event occurs at a point P. The event is seen by two observers in inertial frames S and S', where S' moves with a velocity \mathbf{v} relative to S.

Galilean transformation equations

▲ PITFALL PREVENTION

39.1 The Relationship Between the S and S' Frames

Many of the mathematical representations in this chapter are true *only* for the specified relationship between the S and S' frames. The x and x' axes coincide, except that their origins are different. The y and y' axes (and the z and z' axes), are parallel, but do not coincide due to the displacement of the origin of S' with respect to that of S. We choose the time $t = 0$ to be the instant at which the origins of the two coordinate systems coincide. If the S' frame is moving in the positive x direction relative to S, v is positive; otherwise it is negative.




Figure 39.3 If the velocity of the ether wind relative to the Earth is \mathbf{v} and the velocity of light relative to the ether is \mathbf{c} , then the speed of light relative to the Earth is (a) $c + v$ in the downwind direction, (b) $c - v$ in the upwind direction, and (c) $(c^2 - v^2)^{1/2}$ in the direction perpendicular to the wind.

at rest with respect to the ether. The Galilean velocity transformation equation was expected to hold for observations of light made by an observer in any frame moving at speed v relative to the absolute ether frame. That is, if light travels along the x axis and an observer moves with velocity \mathbf{v} along the x axis, the observer will measure the light to have speed $c \pm v$, depending on the directions of travel of the observer and the light.

Because the existence of a preferred, absolute ether frame would show that light was similar to other classical waves and that Newtonian ideas of an absolute frame were true, considerable importance was attached to establishing the existence of the ether frame. Prior to the late 1800s, experiments involving light traveling in media moving at the highest laboratory speeds attainable at that time were not capable of detecting differences as small as that between c and $c \pm v$. Starting in about 1880, scientists decided to use the Earth as the moving frame in an attempt to improve their chances of detecting these small changes in the speed of light.

As observers fixed on the Earth, we can take the view that we are stationary and that the absolute ether frame containing the medium for light propagation moves past us with speed v . Determining the speed of light under these circumstances is just like determining the speed of an aircraft traveling in a moving air current, or wind; consequently, we speak of an “ether wind” blowing through our apparatus fixed to the Earth.

A direct method for detecting an ether wind would use an apparatus fixed to the Earth to measure the ether wind’s influence on the speed of light. If v is the speed of the ether relative to the Earth, then light should have its maximum speed $c + v$ when propagating downwind, as in Figure 39.3a. Likewise, the speed of light should have its minimum value $c - v$ when the light is propagating upwind, as in Figure 39.3b, and an intermediate value $(c^2 - v^2)^{1/2}$ in the direction perpendicular to the ether wind, as in Figure 39.3c. If the Sun is assumed to be at rest in the ether, then the velocity of the ether wind would be equal to the orbital velocity of the Earth around the Sun, which has a magnitude of approximately 3×10^4 m/s. Because $c = 3 \times 10^8$ m/s, it is necessary to detect a change in speed of about 1 part in 10^4 for measurements in the upwind or downwind directions. However, while such a change is experimentally measurable, all attempts to detect such changes and establish the existence of the ether wind (and hence the absolute frame) proved futile! We explore the classic experimental search for the ether in Section 39.2.

The principle of Galilean relativity refers only to the laws of mechanics. If it is assumed that the laws of electricity and magnetism are the same in all inertial frames, a paradox concerning the speed of light immediately arises. We can understand this by recognizing that Maxwell’s equations seem to imply that the speed of light always has the fixed value 3.00×10^8 m/s in all inertial frames, a result in direct contradiction to what is expected based on the Galilean velocity transformation equation. According to Galilean relativity, the speed of light should *not* be the same in all inertial frames.

To resolve this contradiction in theories, we must conclude that either (1) the laws of electricity and magnetism are not the same in all inertial frames or (2) the Galilean velocity transformation equation is incorrect. If we assume the first alternative, then a preferred reference frame in which the speed of light has the value c must exist and the measured speed must be greater or less than this value in any other reference frame, in accordance with the Galilean velocity transformation equation. If we assume the second alternative, then we are forced to abandon the notions of absolute time and absolute length that form the basis of the Galilean space–time transformation equations.


39.2 The Michelson–Morley Experiment

The most famous experiment designed to detect small changes in the speed of light was first performed in 1881 by Albert A. Michelson (see Section 37.7) and later repeated under various conditions by Michelson and Edward W. Morley (1838–1923). We state at the outset that the outcome of the experiment contradicted the ether hypothesis.

The experiment was designed to determine the velocity of the Earth relative to that of the hypothetical ether. The experimental tool used was the Michelson interferometer, which was discussed in Section 37.7 and is shown again in Figure 39.4. Arm 2 is aligned along the direction of the Earth's motion through space. The Earth moving through the ether at speed v is equivalent to the ether flowing past the Earth in the opposite direction with speed v . This ether wind blowing in the direction opposite the direction of Earth's motion should cause the speed of light measured in the Earth frame to be $c - v$ as the light approaches mirror M_2 and $c + v$ after reflection, where c is the speed of light in the ether frame.

The two light beams reflect from M_1 and M_2 and recombine, and an interference pattern is formed, as discussed in Section 37.7. The interference pattern is observed while the interferometer is rotated through an angle of 90° . This rotation interchanges the speed of the ether wind between the arms of the interferometer. The rotation should cause the fringe pattern to shift slightly but measurably. Measurements failed, however, to show any change in the interference pattern! The Michelson–Morley experiment was repeated at different times of the year when the ether wind was expected to change direction and magnitude, but the results were always the same: **no fringe shift of the magnitude required was ever observed.**²

The negative results of the Michelson–Morley experiment not only contradicted the ether hypothesis but also showed that it was impossible to measure the absolute velocity of the Earth with respect to the ether frame. However, Einstein offered a postulate for his special theory of relativity that places quite a different interpretation on these null results. In later years, when more was known about the nature of light, the idea of an ether that permeates all of space was abandoned. **Light is now understood to be an electromagnetic wave, which requires no medium for its propagation.** As a result, the idea of an ether in which these waves travel became unnecessary.



Active Figure 39.4 According to the ether wind theory, the speed of light should be $c - v$ as the beam approaches mirror M_2 and $c + v$ after reflection.

 **At the Active Figures link at <http://www.pse6.com>, you can adjust the speed of the ether wind to see the effect on the light beams if there were an ether.**

Details of the Michelson–Morley Experiment

To understand the outcome of the Michelson–Morley experiment, let us assume that the two arms of the interferometer in Figure 39.4 are of equal length L . We shall analyze the situation as if there were an ether wind, because that is what Michelson and Morley expected to find. As noted above, the speed of the light beam along arm 2 should be $c - v$ as the beam approaches M_2 and $c + v$ after the beam is reflected. Thus, the time interval for travel to the right is $L/(c - v)$, and the time interval for travel to the left is $L/(c + v)$. The total time interval for the round trip along arm 2 is

$$\Delta t_{\text{arm 2}} = \frac{L}{c + v} + \frac{L}{c - v} = \frac{2Lc}{c^2 - v^2} = \frac{2L}{c} \left(1 - \frac{v^2}{c^2}\right)^{-1}$$

Now consider the light beam traveling along arm 1, perpendicular to the ether wind. Because the speed of the beam relative to the Earth is $(c^2 - v^2)^{1/2}$ in this case (see Fig. 39.3), the time interval for travel for each half of the trip is $L/(c^2 - v^2)^{1/2}$, and the total time interval for the round trip is

$$\Delta t_{\text{arm 1}} = \frac{2L}{(c^2 - v^2)^{1/2}} = \frac{2L}{c} \left(1 - \frac{v^2}{c^2}\right)^{-1/2}$$

Thus, the time difference Δt between the horizontal round trip (arm 2) and the vertical round trip (arm 1) is

$$\Delta t = \Delta t_{\text{arm 2}} - \Delta t_{\text{arm 1}} = \frac{2L}{c} \left[\left(1 - \frac{v^2}{c^2}\right)^{-1} - \left(1 - \frac{v^2}{c^2}\right)^{-1/2} \right]$$

² From an Earth observer's point of view, changes in the Earth's speed and direction of motion in the course of a year are viewed as ether wind shifts. Even if the speed of the Earth with respect to the ether were zero at some time, six months later the speed of the Earth would be 60 km/s with respect to the ether, and as a result a fringe shift should be noticed. No shift has ever been observed, however.

Because $v^2/c^2 \ll 1$, we can simplify this expression by using the following binomial expansion after dropping all terms higher than second order:

$$(1 - x)^n \approx 1 - nx \quad (\text{for } x \ll 1)$$

In our case, $x = v^2/c^2$, and we find that

$$\Delta t = \Delta t_{\text{arm } 2} - \Delta t_{\text{arm } 1} \approx \frac{Lv^2}{c^3} \quad (39.3)$$

This time difference between the two instants at which the reflected beams arrive at the viewing telescope gives rise to a phase difference between the beams, producing an interference pattern when they combine at the position of the telescope. A shift in the interference pattern should be detected when the interferometer is rotated through 90° in a horizontal plane, so that the two beams exchange roles. This rotation results in a time difference twice that given by Equation 39.3. Thus, the path difference that corresponds to this time difference is

$$\Delta d = c(2\Delta t) = \frac{2Lv^2}{c^2}$$

Because a change in path length of one wavelength corresponds to a shift of one fringe, the corresponding fringe shift is equal to this path difference divided by the wavelength of the light:

$$\text{Shift} = \frac{2Lv^2}{\lambda c^2} \quad (39.4)$$

In the experiments by Michelson and Morley, each light beam was reflected by mirrors many times to give an effective path length L of approximately 11 m. Using this value and taking v to be equal to 3.0×10^4 m/s, the speed of the Earth around the Sun, we obtain a path difference of

$$\Delta d = \frac{2(11 \text{ m})(3.0 \times 10^4 \text{ m/s})^2}{(3.0 \times 10^8 \text{ m/s})^2} = 2.2 \times 10^{-7} \text{ m}$$

This extra travel distance should produce a noticeable shift in the fringe pattern. Specifically, using 500-nm light, we expect a fringe shift for rotation through 90° of

$$\text{Shift} = \frac{\Delta d}{\lambda} = \frac{2.2 \times 10^{-7} \text{ m}}{5.0 \times 10^{-7} \text{ m}} \approx 0.44$$

The instrument used by Michelson and Morley could detect shifts as small as 0.01 fringe. However, **it detected no shift whatsoever in the fringe pattern.** Since then, the experiment has been repeated many times by different scientists under a wide variety of conditions, and no fringe shift has ever been detected. Thus, it was concluded that the motion of the Earth with respect to the postulated ether cannot be detected.

Many efforts were made to explain the null results of the Michelson–Morley experiment and to save the ether frame concept and the Galilean velocity transformation equation for light. All proposals resulting from these efforts have been shown to be wrong. No experiment in the history of physics received such valiant efforts to explain the absence of an expected result as did the Michelson–Morley experiment. The stage was set for Einstein, who solved the problem in 1905 with his special theory of relativity.

39.3 Einstein's Principle of Relativity

In the previous section we noted the impossibility of measuring the speed of the ether with respect to the Earth and the failure of the Galilean velocity transformation equation in the case of light. Einstein proposed a theory that boldly removed these

difficulties and at the same time completely altered our notion of space and time.³ He based his special theory of relativity on two postulates:

- 1. The principle of relativity:** The laws of physics must be the same in all inertial reference frames.
- 2. The constancy of the speed of light:** The speed of light in vacuum has the same value, $c = 3.00 \times 10^8$ m/s, in all inertial frames, regardless of the velocity of the observer or the velocity of the source emitting the light.

The first postulate asserts that *all* the laws of physics—those dealing with mechanics, electricity and magnetism, optics, thermodynamics, and so on—are the same in all reference frames moving with constant velocity relative to one another. This postulate is a sweeping generalization of the principle of Galilean relativity, which refers only to the laws of mechanics. From an experimental point of view, Einstein’s principle of relativity means that any kind of experiment (measuring the speed of light, for example) performed in a laboratory at rest must give the same result when performed in a laboratory moving at a constant velocity with respect to the first one. Hence, no preferred inertial reference frame exists, and it is impossible to detect absolute motion.

Note that postulate 2 is required by postulate 1: if the speed of light were not the same in all inertial frames, measurements of different speeds would make it possible to distinguish between inertial frames; as a result, a preferred, absolute frame could be identified, in contradiction to postulate 1.


Although the Michelson–Morley experiment was performed before Einstein published his work on relativity, it is not clear whether or not Einstein was aware of the details of the experiment. Nonetheless, the null result of the experiment can be readily understood within the framework of Einstein’s theory. According to his principle of relativity, the premises of the Michelson–Morley experiment were incorrect. In the process of trying to explain the expected results, we stated that when light traveled against the ether wind its speed was $c - v$, in accordance with the Galilean velocity transformation equation. However, if the state of motion of the observer or of the source has no influence on the value found for the speed of light, one always measures the value to be c . Likewise, the light makes the return trip after reflection from the mirror at speed c , not at speed $c + v$. Thus, the motion of the Earth does not influence the fringe pattern observed in the Michelson–Morley experiment, and a null result should be expected.

If we accept Einstein’s theory of relativity, we must conclude that relative motion is unimportant when measuring the speed of light. At the same time, we shall see that we must alter our common-sense notion of space and time and be prepared for some surprising consequences. It may help as you read the pages ahead to keep in mind that our common-sense ideas are based on a lifetime of everyday experiences and not on observations of objects moving at hundreds of thousands of kilometers per second. Thus, these results will seem strange, but that is only because we have no experience with them.

39.4 Consequences of the Special Theory of Relativity

Before we discuss the consequences of Einstein’s special theory of relativity, we must first understand how an observer located in an inertial reference frame describes an event. As mentioned earlier, an event is an occurrence describable by three space

³ A. Einstein, “On the Electrodynamics of Moving Bodies,” *Ann. Physik* 17:891, 1905. For an English translation of this article and other publications by Einstein, see the book by H. Lorentz, A. Einstein, H. Minkowski, and H. Weyl, *The Principle of Relativity*, Dover, 1958.



Albert Einstein
German-American Physicist
(1879–1955)

Einstein, one of the greatest physicists of all times, was born in Ulm, Germany. In 1905, at the age of 26, he published four scientific papers that revolutionized physics. Two of these papers were concerned with what is now considered his most important contribution: the special theory of relativity.

In 1916, Einstein published his work on the general theory of relativity. The most dramatic prediction of this theory is the degree to which light is deflected by a gravitational field. Measurements made by astronomers on bright stars in the vicinity of the eclipsed Sun in 1919 confirmed Einstein’s prediction, and as a result Einstein became a world celebrity.

Einstein was deeply disturbed by the development of quantum mechanics in the 1920s despite his own role as a scientific revolutionary. In particular, he could never accept the probabilistic view of events in nature that is a central feature of quantum theory. The last few decades of his life were devoted to an unsuccessful search for a unified theory that would combine gravitation and electromagnetism. (*AIP Niels Bohr Library*)

coordinates and one time coordinate. Observers in different inertial frames will describe the same event with coordinates that have different values.

As we examine some of the consequences of relativity in the remainder of this section, we restrict our discussion to the concepts of simultaneity, time intervals, and lengths, all three of which are quite different in relativistic mechanics from what they are in Newtonian mechanics. For example, in relativistic mechanics the distance between two points and the time interval between two events depend on the frame of reference in which they are measured. That is, **in relativistic mechanics there is no such thing as an absolute length or absolute time interval.** Furthermore, **events at different locations that are observed to occur simultaneously in one frame are not necessarily observed to be simultaneous in another frame moving uniformly with respect to the first.**

Simultaneity and the Relativity of Time

A basic premise of Newtonian mechanics is that a universal time scale exists that is the same for all observers. In fact, Newton wrote that “Absolute, true, and mathematical time, of itself, and from its own nature, flows equably without relation to anything external.” Thus, Newton and his followers simply took simultaneity for granted. In his special theory of relativity, Einstein abandoned this assumption.

Einstein devised the following thought experiment to illustrate this point. A boxcar moves with uniform velocity, and two lightning bolts strike its ends, as illustrated in Figure 39.5a, leaving marks on the boxcar and on the ground. The marks on the boxcar are labeled A' and B' , and those on the ground are labeled A and B . An observer O' moving with the boxcar is midway between A' and B' , and a ground observer O is midway between A and B . The events recorded by the observers are the striking of the boxcar by the two lightning bolts.

The light signals emitted from A and B at the instant at which the two bolts strike reach observer O at the same time, as indicated in Figure 39.5b. This observer realizes that the signals have traveled at the same speed over equal distances, and so rightly concludes that the events at A and B occurred simultaneously. Now consider the same events as viewed by observer O' . By the time the signals have reached observer O , observer O' has moved as indicated in Figure 39.5b. Thus, the signal from B' has already swept past O' , but the signal from A' has not yet reached O' . In other words, O' sees the signal from B' before seeing the signal from A' . According to Einstein, *the two observers must find that light travels at the same speed.* Therefore, observer O' concludes that the lightning strikes the front of the boxcar before it strikes the back.

This thought experiment clearly demonstrates that the two events that appear to be simultaneous to observer O do not appear to be simultaneous to observer O' .

▲ PITFALL PREVENTION

39.2 Who's Right?

You might wonder which observer in Fig. 39.5 is correct concerning the two lightning strikes. *Both are correct*, because the principle of relativity states that *there is no preferred inertial frame of reference.* Although the two observers reach different conclusions, both are correct in their own reference frame because the concept of simultaneity is not absolute. This, in fact, is the central point of relativity—any uniformly moving frame of reference can be used to describe events and do physics.





Figure 39.5 (a) Two lightning bolts strike the ends of a moving boxcar. (b) The events appear to be simultaneous to the stationary observer O , standing midway between A and B . The events do not appear to be simultaneous to observer O' , who claims that the front of the car is struck before the rear. Note that in (b) the leftward-traveling light signal has already passed O' but the rightward-traveling signal has not yet reached O' .



Active Figure 39.6 (a) A mirror is fixed to a moving vehicle, and a light pulse is sent out by observer O' at rest in the vehicle. (b) Relative to a stationary observer O standing alongside the vehicle, the mirror and O' move with a speed v . Note that what observer O measures for the distance the pulse travels is greater than $2d$. (c) The right triangle for calculating the relationship between Δt and Δt_p .

 **At the Active Figures link at <http://www.pse6.com>, you can observe the bouncing of the light pulse for various speeds of the train.**

In other words,

two events that are simultaneous in one reference frame are in general not simultaneous in a second frame moving relative to the first. That is, simultaneity is not an absolute concept but rather one that depends on the state of motion of the observer.

Einstein's thought experiment demonstrates that two observers can disagree on the simultaneity of two events. **This disagreement, however, depends on the transit time of light to the observers and, therefore, does not demonstrate the deeper meaning of relativity.** In relativistic analyses of high-speed situations, relativity shows that simultaneity is relative even when the transit time is subtracted out. In fact, all of the relativistic effects that we will discuss from here on will assume that we are ignoring differences caused by the transit time of light to the observers.

Time Dilation

We can illustrate the fact that observers in different inertial frames can measure different time intervals between a pair of events by considering a vehicle moving to the right with a speed v , such as the boxcar shown in Figure 39.6a. A mirror is fixed to the ceiling of the vehicle, and observer O' at rest in the frame attached to the vehicle holds a flashlight a distance d below the mirror. At some instant, the flashlight emits a pulse of light directed toward the mirror (event 1), and at some later time after reflecting from the mirror, the pulse arrives back at the flashlight (event 2). Observer O' carries a clock and uses it to measure the time interval Δt_p between these two events. (The subscript p stands for *proper*, as we shall see in a moment.) Because the light pulse has a speed c , the time interval required for the pulse to travel from O' to the mirror and back is

$$\Delta t_p = \frac{\text{distance traveled}}{\text{speed}} = \frac{2d}{c} \quad (39.5)$$

Now consider the same pair of events as viewed by observer O in a second frame, as shown in Figure 39.6b. According to this observer, the mirror and flashlight are moving to the right with a speed v , and as a result the sequence of events appears entirely different. By the time the light from the flashlight reaches the mirror, the mirror has moved to the right a distance $v \Delta t/2$, where Δt is the time interval required for the light to travel from O' to the mirror and back to O' as measured by O . In other words, O concludes that, because of the motion of the vehicle, if the light is to hit the mirror, it must leave the

flashlight at an angle with respect to the vertical direction. Comparing Figure 39.6a and b, we see that the light must travel farther in (b) than in (a). (Note that neither observer “knows” that he or she is moving. Each is at rest in his or her own inertial frame.)

According to the second postulate of the special theory of relativity, both observers must measure c for the speed of light. Because the light travels farther according to O , it follows that the time interval Δt measured by O is longer than the time interval Δt_p measured by O' . To obtain a relationship between these two time intervals, it is convenient to use the right triangle shown in Figure 39.6c. The Pythagorean theorem gives

$$\left(\frac{c \Delta t}{2}\right)^2 = \left(\frac{v \Delta t}{2}\right)^2 + d^2$$

Solving for Δt gives

$$\Delta t = \frac{2d}{\sqrt{c^2 - v^2}} = \frac{2d}{c \sqrt{1 - \frac{v^2}{c^2}}} \tag{39.6}$$

Because $\Delta t_p = 2d/c$, we can express this result as

$$\Delta t = \frac{\Delta t_p}{\sqrt{1 - \frac{v^2}{c^2}}} = \gamma \Delta t_p \tag{39.7}$$

where

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \tag{39.8}$$

Because γ is always greater than unity, this result says that **the time interval Δt measured by an observer moving with respect to a clock is longer than the time interval Δt_p measured by an observer at rest with respect to the clock.** This effect is known as **time dilation**.

We can see that time dilation is not observed in our everyday lives by considering the factor γ . This factor deviates significantly from a value of 1 only for very high speeds, as shown in Figure 39.7 and Table 39.1. For example, for a speed of $0.1c$, the value of γ is 1.005. Thus, there is a time dilation of only 0.5% at one-tenth the speed of light. Speeds that we encounter on an everyday basis are far slower than this, so we do not see time dilation in normal situations.

The time interval Δt_p in Equations 39.5 and 39.7 is called the **proper time interval**. (In German, Einstein used the term *Eigenzeit*, which means “own-time.”) In

Time dilation

Table 39.1

Approximate Values for γ at Various Speeds	
v/c	γ
0.001 0	1.000 000 5
0.010	1.000 05
0.10	1.005
0.20	1.021
0.30	1.048
0.40	1.091
0.50	1.155
0.60	1.250
0.70	1.400
0.80	1.667
0.90	2.294
0.92	2.552
0.94	2.931
0.96	3.571
0.98	5.025
0.99	7.089
0.995	10.01
0.999	22.37




Figure 39.7 Graph of γ versus v . As the speed approaches that of light, γ increases rapidly.

general, **the proper time interval is the time interval between two events measured by an observer who sees the events occur at the same point in space.**

If a clock is moving with respect to you, the time interval between ticks of the moving clock is observed to be longer than the time interval between ticks of an identical clock in your reference frame. Thus, it is often said that a moving clock is measured to run more slowly than a clock in your reference frame by a factor γ . This is true for mechanical clocks as well as for the light clock just described. We can generalize this result by stating that all physical processes, including chemical and biological ones, are measured to slow down when those processes occur in a frame moving with respect to the observer. For example, the heartbeat of an astronaut moving through space would keep time with a clock inside the spacecraft. Both the astronaut's clock and heartbeat would be measured to slow down according to an observer on Earth comparing time intervals with his own clock (although the astronaut would have no sensation of life slowing down in the spacecraft).

Quick Quiz 39.3 Suppose the observer O' on the train in Figure 39.6 aims her flashlight at the far wall of the boxcar and turns it on and off, sending a pulse of light toward the far wall. Both O' and O measure the time interval between when the pulse leaves the flashlight and it hits the far wall. Which observer measures the proper time interval between these two events? (a) O' (b) O (c) both observers (d) neither observer.

Quick Quiz 39.4 A crew watches a movie that is two hours long in a spacecraft that is moving at high speed through space. Will an Earthbound observer, who is watching the movie through a powerful telescope, measure the duration of the movie to be (a) longer than, (b) shorter than, or (c) equal to two hours?

Strange as it may seem, time dilation is a verifiable phenomenon. An experiment reported by Hafele and Keating provided direct evidence of time dilation.⁴ Time intervals measured with four cesium atomic clocks in jet flight were compared with time intervals measured by Earth-based reference atomic clocks. In order to compare these results with theory, many factors had to be considered, including periods of speeding up and slowing down relative to the Earth, variations in direction of travel, and the fact that the gravitational field experienced by the flying clocks was weaker than that experienced by the Earth-based clock. The results were in good agreement with the predictions of the special theory of relativity and can be explained in terms of the relative motion between the Earth and the jet aircraft. In their paper, Hafele and Keating stated that “Relative to the atomic time scale of the U.S. Naval Observatory, the flying clocks lost 59 ± 10 ns during the eastward trip and gained 273 ± 7 ns during the westward trip. . . . These results provide an unambiguous empirical resolution of the famous clock paradox with macroscopic clocks.”

Another interesting example of time dilation involves the observation of *muons*, unstable elementary particles that have a charge equal to that of the electron and a mass 207 times that of the electron. (We will study the muon and other particles in Chapter 46.) Muons can be produced by the collision of cosmic radiation with atoms high in the atmosphere. Slow-moving muons in the laboratory have a lifetime which is measured to be the proper time interval $\Delta t_p = 2.2 \mu\text{s}$. If we assume that the speed of atmospheric muons is close to the speed of light, we find that these particles can travel a distance of approximately $(3.0 \times 10^8 \text{ m/s})(2.2 \times 10^{-6} \text{ s}) \approx 6.6 \times 10^2 \text{ m}$ before they decay (Fig. 39.8a). Hence, they are unlikely to reach the surface of the Earth from

PITFALL PREVENTION

39.3 The Proper Time Interval

It is *very* important in relativistic calculations to correctly identify the observer who measures the proper time interval. The proper time interval between two events is always the time interval measured by an observer for whom the two events take place at the same position.




Figure 39.8 (a) Without relativistic considerations, muons created in the atmosphere and traveling downward with a speed of $0.99c$ travel only about $6.6 \times 10^2 \text{ m}$ before decaying with an average lifetime of $2.2 \mu\text{s}$. Thus, very few muons reach the surface of the Earth. (b) With relativistic considerations, the muon's lifetime is dilated according to an observer on Earth. As a result, according to this observer, the muon can travel about $4.8 \times 10^3 \text{ m}$ before decaying. This results in many of them arriving at the surface.

⁴ J. C. Hafele and R. E. Keating, “Around the World Atomic Clocks: Relativistic Time Gains Observed,” *Science*, 177:168, 1972.




Figure 39.9 Decay curves for muons at rest and for muons traveling at a speed of $0.9994c$.

high in the atmosphere where they are produced. However, experiments show that a large number of muons *do* reach the surface. The phenomenon of time dilation explains this effect. As measured by an observer on Earth, the muons have a dilated lifetime equal to $\gamma \Delta t_p$. For example, for $v = 0.99c$, $\gamma \approx 7.1$ and $\gamma \Delta t_p \approx 16 \mu\text{s}$. Hence, the average distance traveled by the muons in this time as measured by an observer on Earth is approximately $(0.99)(3.0 \times 10^8 \text{ m/s})(16 \times 10^{-6} \text{ s}) \approx 4.8 \times 10^3 \text{ m}$, as indicated in Figure 39.8b.

In 1976, at the laboratory of the European Council for Nuclear Research (CERN) in Geneva, muons injected into a large storage ring reached speeds of approximately $0.9994c$. Electrons produced by the decaying muons were detected by counters around the ring, enabling scientists to measure the decay rate and hence the muon lifetime. The lifetime of the moving muons was measured to be approximately 30 times as long as that of the stationary muon (Fig. 39.9), in agreement with the prediction of relativity to within two parts in a thousand.

Example 39.1 What Is the Period of the Pendulum?

The period of a pendulum is measured to be 3.00 s in the reference frame of the pendulum. What is the period when measured by an observer moving at a speed of $0.950c$ relative to the pendulum?

Solution To conceptualize this problem, let us change frames of reference. Instead of the observer moving at $0.950c$, we can take the equivalent point of view that the observer is at rest and the pendulum is moving at $0.950c$ past the stationary observer. Hence, the pendulum is an example of a clock moving at high speed with respect to an observer and we can categorize this problem as one involving time dilation.

To analyze the problem, note that the proper time interval, measured in the rest frame of the pendulum, is $\Delta t_p = 3.00 \text{ s}$. Because a clock moving with respect to an observer is measured to run more slowly than a stationary clock by a factor γ , Equation 39.7 gives

$$\begin{aligned} \Delta t &= \gamma \Delta t_p = \frac{1}{\sqrt{1 - \frac{(0.950c)^2}{c^2}}} \Delta t_p = \frac{1}{\sqrt{1 - 0.902}} \Delta t_p \\ &= (3.20)(3.00 \text{ s}) = 9.60 \text{ s} \end{aligned}$$

To finalize this problem, we see that indeed a moving pendulum is measured to take longer to complete a period

than a pendulum at rest does. The period increases by a factor of $\gamma = 3.20$. We see that this is consistent with Table 39.1, where this value lies between those for γ for $v/c = 0.94$ and $v/c = 0.96$.

What If? What if we increase the speed of the observer by 5.00%? Does the dilated time interval increase by 5.00%?

Answer Based on the highly nonlinear behavior of γ as a function of v in Figure 39.7, we would guess that the increase in Δt would be different from 5.00%. Increasing v by 5.00% gives us

$$v_{\text{new}} = (1.0500)(0.950c) = 0.9975c$$

(Because γ varies so rapidly with v when v is this large, we will keep one additional significant figure until the final answer.) If we perform the time dilation calculation again, we find that

$$\begin{aligned} \Delta t &= \gamma \Delta t_p = \frac{1}{\sqrt{1 - \frac{(0.9975c)^2}{c^2}}} \Delta t_p = \frac{1}{\sqrt{1 - 0.9950}} \Delta t_p \\ &= (14.15)(3.00 \text{ s}) = 42.5 \text{ s} \end{aligned}$$

Thus, the 5.00% increase in speed has caused over a 300% increase in the dilated time!

Example 39.2 How Long Was Your Trip?

Suppose you are driving your car on a business trip and are traveling at 30 m/s. Your boss, who is waiting at your destination, expects the trip to take 5.0 h. When you arrive late, your excuse is that your car clock registered the passage of 5.0 h but that you were driving fast and so your clock ran more slowly than your boss's clock. If your car clock actually did indicate a 5.0-h trip, how much time passed on your boss's clock, which was at rest on the Earth?

Solution We begin by calculating γ from Equation 39.8:

$$\begin{aligned}\gamma &= \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{1}{\sqrt{1 - \frac{(3 \times 10^1 \text{ m/s})^2}{(3 \times 10^8 \text{ m/s})^2}}} \\ &= \frac{1}{\sqrt{1 - 10^{-14}}}\end{aligned}$$

If you try to determine this value on your calculator, you will probably obtain $\gamma = 1$. However, if we perform a binomial expansion, we can more precisely determine the value as

$$\gamma = (1 - 10^{-14})^{-1/2} \approx 1 + \frac{1}{2}(10^{-14}) = 1 + 5.0 \times 10^{-15}$$

This result indicates that at typical automobile speeds, γ is not much different from 1.

Applying Equation 39.7, we find Δt , the time interval measured by your boss, to be

$$\begin{aligned}\Delta t &= \gamma \Delta t_p = (1 + 5.0 \times 10^{-15})(5.0 \text{ h}) \\ &= 5.0 \text{ h} + 2.5 \times 10^{-14} \text{ h} = \mathbf{5.0 \text{ h} + 0.09 \text{ ns}}\end{aligned}$$

Your boss's clock would be only 0.09 ns ahead of your car clock. You might want to think of another excuse!

The Twin Paradox

An intriguing consequence of time dilation is the so-called *twin paradox* (Fig. 39.10). Consider an experiment involving a set of twins named Speedo and Goslo. When they are 20 yr old, Speedo, the more adventuresome of the two, sets out on an epic journey to Planet X, located 20 ly from the Earth. (Note that 1 lightyear (ly) is the distance light travels through free space in 1 year.) Furthermore, Speedo's spacecraft is capable of reaching a speed of $0.95c$ relative to the inertial frame of his twin brother back home. After reaching Planet X, Speedo becomes homesick and immediately returns to the Earth at the same speed $0.95c$. Upon his return, Speedo is shocked to discover that Goslo has aged 42 yr and is now 62 yr old. Speedo, on the other hand, has aged only 13 yr.

At this point, it is fair to raise the following question—which twin is the traveler and which is really younger as a result of this experiment? From Goslo's frame of reference, he was at rest while his brother traveled at a high speed away from him and then came back. According to Speedo, however, he himself remained stationary while Goslo and the Earth raced away from him and then headed back. This leads to an apparent




Figure 39.10 (a) As one twin leaves his brother on the Earth, both are the same age. (b) When Speedo returns from his journey to Planet X, he is younger than his twin Goslo.

contradiction due to the apparent symmetry of the observations. Which twin has developed signs of excess aging?

The situation in our current problem is actually not symmetrical. To resolve this apparent paradox, recall that the special theory of relativity describes observations made in inertial frames of reference moving relative to each other. Speedo, the space traveler, must experience a series of accelerations during his journey because he must fire his rocket engines to slow down and start moving back toward Earth. As a result, his speed is not always uniform, and consequently he is not in an inertial frame. Therefore, there is no paradox—only Goslo, who is always in a single inertial frame, can make correct predictions based on special relativity. During each passing year noted by Goslo, slightly less than 4 months elapses for Speedo.

Only Goslo, who is in a single inertial frame, can apply the simple time-dilation formula to Speedo's trip. Thus, Goslo finds that instead of aging 42 yr, Speedo ages only $(1 - v^2/c^2)^{1/2}(42 \text{ yr}) = 13 \text{ yr}$. Thus, according to Goslo, Speedo spends 6.5 yr traveling to Planet X and 6.5 yr returning, for a total travel time of 13 yr, in agreement with our earlier statement.

Quick Quiz 39.5 Suppose astronauts are paid according to the amount of time they spend traveling in space. After a long voyage traveling at a speed approaching c , would a crew rather be paid according to (a) an Earth-based clock, (b) their spacecraft's clock, or (c) either clock?

▲ PITFALL PREVENTION

39.4 The Proper Length

As with the proper time interval, it is *very* important in relativistic calculations to correctly identify the observer who measures the proper length. The proper length between two points in space is always the length measured by an observer at rest with respect to the points. Often the proper time interval and the proper length are *not* measured by the same observer.

Length Contraction

The measured distance between two points also depends on the frame of reference. **The proper length L_p of an object is the length measured by someone at rest relative to the object.** The length of an object measured by someone in a reference frame that is moving with respect to the object is always less than the proper length. This effect is known as **length contraction**.

Consider a spacecraft traveling with a speed v from one star to another. There are two observers: one on the Earth and the other in the spacecraft. The observer at rest on the Earth (and also assumed to be at rest with respect to the two stars) measures the distance between the stars to be the proper length L_p . According to this observer, the time interval required for the spacecraft to complete the voyage is $\Delta t = L_p/v$. The passages of the two stars by the spacecraft occur at the same position for the space traveler. Thus, the space traveler measures the proper time interval Δt_p . Because of time dilation, the proper time interval is related to the Earth-measured time interval by $\Delta t_p = \Delta t/\gamma$. Because the space traveler reaches the second star in the time Δt_p , he or she concludes that the distance L between the stars is

$$L = v \Delta t_p = v \frac{\Delta t}{\gamma}$$

Because the proper length is $L_p = v \Delta t$, we see that


$$L = \frac{L_p}{\gamma} = L_p \sqrt{1 - \frac{v^2}{c^2}} \quad (39.9)$$

Length contraction

where $\sqrt{1 - v^2/c^2}$ is a factor less than unity. **If an object has a proper length L_p when it is measured by an observer at rest with respect to the object, then when it moves with speed v in a direction parallel to its length, its length L is measured to be shorter according to $L = L_p \sqrt{1 - v^2/c^2} = L_p/\gamma$.**

For example, suppose that a meter stick moves past a stationary Earth observer with speed v , as in Figure 39.11. The length of the stick as measured by an observer in a frame attached to the stick is the proper length L_p shown in Figure 39.11a. The length of the stick L measured by the Earth observer is shorter than L_p by the factor $(1 - v^2/c^2)^{1/2}$. Note that **length contraction takes place only along the direction of motion**.

The proper length and the proper time interval are defined differently. The proper length is measured by an observer for whom the end points of the length remain fixed in space. The proper time interval is measured by someone for whom the two events take place at the same position in space. As an example of this point, let us return to the decaying muons moving at speeds close to the speed of light. An observer in the muon's reference frame would measure the proper lifetime, while an Earth-based observer would measure the proper length (the distance from creation to decay in Figure 39.8). In the muon's reference frame, there is no time dilation but the distance of travel to the surface is observed to be shorter when measured in this frame. Likewise, in the Earth observer's reference frame, there is time dilation, but the distance of travel is measured to be the proper length. Thus, when calculations on the muon are performed in both frames, the outcome of the experiment in one frame is the same as the outcome in the other frame—more muons reach the surface than would be predicted without relativistic effects.



Active Figure 39.11 (a) A meter stick measured by an observer in a frame attached to the stick (that is, both have the same velocity) has its proper length L_p . (b) The stick measured by an observer in a frame in which the stick has a velocity \mathbf{v} relative to the frame is measured to be shorter than its proper length L_p by a factor $(1 - v^2/c^2)^{1/2}$.

 **At the Active Figures link at <http://www.pse6.com>, you can view the meter stick from the points of view of two observers to compare the measured length of the stick.**

Quick Quiz 39.6 You are packing for a trip to another star. During the journey, you will be traveling at $0.99c$. You are trying to decide whether you should buy smaller sizes of your clothing, because you will be thinner on your trip, due to length contraction. Also, you are considering saving money by reserving a smaller cabin to sleep in, because you will be shorter when you lie down. Should you (a) buy smaller sizes of clothing, (b) reserve a smaller cabin, (c) do neither of these, or (d) do both of these?

Quick Quiz 39.7 You are observing a spacecraft moving away from you. You measure it to be shorter than when it was at rest on the ground next to you. You also see a clock through the spacecraft window, and you observe that the passage of time on the clock is measured to be slower than that of the watch on your wrist. Compared to when the spacecraft was on the ground, what do you measure if the spacecraft turns around and comes *toward* you at the same speed? (a) The spacecraft is measured to be longer and the clock runs faster. (b) The spacecraft is measured to be longer and the clock runs slower. (c) The spacecraft is measured to be shorter and the clock runs faster. (d) The spacecraft is measured to be shorter and the clock runs slower.

Space–Time Graphs

It is sometimes helpful to make a *space–time graph*, in which ct is the ordinate and position x is the abscissa. The twin paradox is displayed in such a graph in Figure 39.12




Figure 39.12 The twin paradox on a space–time graph. The twin who stays on the Earth has a world-line along the ct axis. The path of the traveling twin through space–time is represented by a world-line that changes direction.

from the point of view of Goslo. A path through space–time is called a **world-line**. At the origin, the world-lines of Speedo and Goslo coincide because the twins are in the same location at the same time. After Speedo leaves on his trip, his world-line diverges from that of his brother. Goslo’s world-line is vertical because he remains fixed in location. At their reunion, the two world-lines again come together. Note that it would be impossible for Speedo to have a world-line that crossed the path of a light beam that left the Earth when he did. To do so would require him to have a speed greater than c (not possible, as shown in Sections 39.6 and 39.7).

World-lines for light beams are diagonal lines on space–time graphs, typically drawn at 45° to the right or left of vertical (assuming that the x and ct axes have the same scales), depending on whether the light beam is traveling in the direction of increasing or decreasing x . These two world-lines mean that all possible future events for Goslo and Speedo lie within two 45° lines extending from the origin. Either twin’s presence at an event outside this “light cone” would require that twin to move at a speed greater than c , which we have said is not possible. Also, the only past events that Goslo and Speedo could have experienced occurred within two similar 45° world-lines that approach the origin from below the x axis.

Example 39.3 The Contraction of a Spacecraft

A spacecraft is measured to be 120.0 m long and 20.0 m in diameter while at rest relative to an observer. If this spacecraft now flies by the observer with a speed of $0.99c$, what length and diameter does the observer measure?

Solution From Equation 39.9, the length measured by the

observer is

$$L = L_p \sqrt{1 - \frac{v^2}{c^2}} = (120.0 \text{ m}) \sqrt{1 - \frac{(0.99c)^2}{c^2}} = 17 \text{ m}$$

The diameter measured by the observer is still 20.0 m because the diameter is a dimension perpendicular to the motion and length contraction occurs only along the direction of motion.

Example 39.4 The Pole-in-the-Barn Paradox

Interactive

The twin paradox, discussed earlier, is a classic “paradox” in relativity. Another classic “paradox” is this: Suppose a runner moving at $0.75c$ carries a horizontal pole 15 m long toward a barn that is 10 m long. The barn has front and rear doors. An observer on the ground can instantly and simultaneously open and close the two doors by remote control. When the runner and the pole are inside the barn, the ground observer closes and then opens both doors so that the runner and pole are momentarily captured inside the barn and then proceed to exit the barn from the back door. Do both the runner and the ground observer agree that the runner makes it safely through the barn?

Solution From our everyday experience, we would be surprised to see a 15-m pole fit inside a 10-m barn. But the pole is in motion with respect to the ground observer, who measures the pole to be contracted to a length L_{pole} , where

$$L_{\text{pole}} = L_p \sqrt{1 - \frac{v^2}{c^2}} = (15 \text{ m}) \sqrt{1 - (0.75)^2} = 9.9 \text{ m}$$

Thus, the ground observer measures the pole to be slightly shorter than the barn and there is no problem with momentarily capturing the pole inside it. The “paradox” arises when we consider the runner’s point of view. The runner

sees the barn contracted to

$$L_{\text{barn}} = L_p \sqrt{1 - \frac{v^2}{c^2}} = (10 \text{ m}) \sqrt{1 - (0.75)^2} = 6.6 \text{ m}$$

Because the pole is in the rest frame of the runner, the runner measures it to have its proper length of 15 m. How can a 15-m pole fit inside a 6.6-m barn? While this is the classic question that is often asked, this is not the question we have asked, because it is not the important question. We asked *if the runner can make it safely through the barn*.

The resolution of the “paradox” lies in the relativity of simultaneity. The closing of the two doors is measured to be simultaneous by the ground observer. Because the doors are at different positions, however, they do not close simultaneously as measured by the runner. The rear door closes and then opens first, allowing the leading edge of the pole to exit. The front door of the barn does not close until the trailing edge of the pole passes by.

We can analyze this using a space–time graph. Figure 39.13a is a space–time graph from the ground observer’s point of view. We choose $x = 0$ as the position of the front door of the barn and $t = 0$ as the instant at which the leading end of the pole is located at the front door of the barn. The world-lines for the two ends of the barn are separated by 10 m and are vertical because the barn is not moving relative to this observer. For the pole, we follow two tilted world-lines, one




Figure 39.13 (Example 39.4) Space–time graphs for the pole-in-the-barn paradox. (a) From the ground observer’s point of view, the world-lines for the front and back doors of the barn are vertical lines. The world-lines for the ends of the pole are tilted and are 9.9 m apart horizontally. The front door of the barn is at $x = 0$, and the leading end of the pole enters the front door at $t = 0$. The entire pole is inside the barn at the time indicated by the dashed line. (b) From the runner’s point of view, the world-lines for the ends of the pole are vertical. The barn is moving in the negative direction, so the world-lines for the front and back doors are tilted to the left. The leading end of the pole exits the back door before the trailing end arrives at the front door.

for each end of the moving pole. These world-lines are 9.9 m apart horizontally, which is the contracted length seen by the ground observer. As seen in Figure 39.13a, at one instant, the pole is entirely within the barn.

Figure 39.13b shows the space–time graph according to the runner. Here, the world-lines for the pole are separated by 15 m and are vertical because the pole is at rest in the runner’s frame of reference. The barn is hurtling *toward* the runner, so the world-lines for the front and rear doors of the barn are tilted in the opposite direction compared to Figure 39.13a. The world-lines for the front and rear doors are separated by 6.6 m, the contracted length as seen by the runner. Notice that the front of the pole leaves the rear door of the barn long before the back of the pole enters the barn. Thus, the opening of the rear door occurs before the closing of the front door.

From the ground observer’s point of view, the time at which the trailing end of the pole enters the barn is found from

$$\Delta t = t - 0 = t = \frac{\Delta x}{v} = \frac{9.9 \text{ m}}{0.75c} = \frac{13.2 \text{ m}}{c}$$

Thus, the pole should be completely inside the barn at a time corresponding to $ct = 13.2 \text{ m}$. This is consistent with the point on the ct axis in Figure 39.13a where the pole is inside the barn.

From the runner’s point of view, the time at which the leading end of the pole leaves the barn is found from

$$\Delta t = t - 0 = t = \frac{\Delta x}{v} = \frac{6.6 \text{ m}}{0.75c} = \frac{8.8 \text{ m}}{c}$$

leading to $ct = 8.8 \text{ m}$. This is consistent with the point on the ct axis in Figure 39.13b where the back door of the barn arrives at the leading end of the pole. Finally, the time at which the trailing end of the pole enters the front door of the barn is found from

$$\Delta t = t - 0 = t = \frac{\Delta x}{v} = \frac{15 \text{ m}}{0.75c} = \frac{20 \text{ m}}{c}$$

This gives $ct = 20 \text{ m}$, which agrees with the instant shown in Figure 39.13b.



Investigate the pole-in-the-barn paradox at the Interactive Worked Example link at <http://www.pse6.com>.

Example 39.5 A Voyage to Sirius

An astronaut takes a trip to Sirius, which is located a distance of 8 lightyears from the Earth. The astronaut measures the time of the one-way journey to be 6 yr. If the spaceship moves at a constant speed of $0.8c$, how can the 8-ly distance be reconciled with the 6-yr trip time measured by the astronaut?

Solution The distance of 8 ly represents the proper length from the Earth to Sirius measured by an observer seeing both objects nearly at rest. The astronaut sees Sirius approaching her at $0.8c$ but also sees the distance

contracted to

$$\frac{8 \text{ ly}}{\gamma} = (8 \text{ ly}) \sqrt{1 - \frac{v^2}{c^2}} = (8 \text{ ly}) \sqrt{1 - \frac{(0.8c)^2}{c^2}} = 5 \text{ ly}$$

Thus, the travel time measured on her clock is

$$\Delta t = \frac{d}{v} = \frac{5 \text{ ly}}{0.8c} = 6 \text{ yr}$$

Note that we have used the value for the speed of light as $c = 1 \text{ ly/yr}$.

What If? What if this trip is observed with a very powerful telescope by a technician in Mission Control on Earth? At what time will this technician see that the astronaut has arrived at Sirius?

Answer The time interval that the technician will measure for the astronaut to arrive is

$$\Delta t = \frac{d}{v} = \frac{8 \text{ ly}}{0.8c} = 10 \text{ yr}$$

In order for the technician to see the arrival, the light from the scene of the arrival must travel back to Earth and enter

the telescope. This will require a time interval of

$$\Delta t = \frac{d}{v} = \frac{8 \text{ ly}}{c} = 8 \text{ yr}$$

Thus, the technician sees the arrival after 10 yr + 8 yr = 18 yr. Notice that if the astronaut immediately turns around and comes back home, she arrives, according to the technician, 20 years after leaving, only 2 years after he saw her arrive! In addition, she would have aged by only 12 years.

The Relativistic Doppler Effect

Another important consequence of time dilation is the shift in frequency found for light emitted by atoms in motion as opposed to light emitted by atoms at rest. This phenomenon, known as the Doppler effect, was introduced in Chapter 17 as it pertains to sound waves. In the case of sound, the motion of the source with respect to the medium of propagation can be distinguished from the motion of the observer with respect to the medium. Light waves must be analyzed differently, however, because they require no medium of propagation, and no method exists for distinguishing the motion of a light source from the motion of the observer.

If a light source and an observer approach each other with a relative speed v , the frequency f_{obs} measured by the observer is

$$f_{\text{obs}} = \frac{\sqrt{1 + v/c}}{\sqrt{1 - v/c}} f_{\text{source}} \tag{39.10}$$

where f_{source} is the frequency of the source measured in its rest frame. Note that this relativistic Doppler shift equation, unlike the Doppler shift equation for sound, depends only on the relative speed v of the source and observer and holds for relative speeds as great as c . As you might expect, the equation predicts that $f_{\text{obs}} > f_{\text{source}}$ when the source and observer approach each other. We obtain the expression for the case in which the source and observer recede from each other by substituting negative values for v in Equation 39.10.

The most spectacular and dramatic use of the relativistic Doppler effect is the measurement of shifts in the frequency of light emitted by a moving astronomical object such as a galaxy. Light emitted by atoms and normally found in the extreme violet region of the spectrum is shifted toward the red end of the spectrum for atoms in other galaxies—indicating that these galaxies are *receding* from us. The American astronomer Edwin Hubble (1889–1953) performed extensive measurements of this *red shift* to confirm that most galaxies are moving away from us, indicating that the Universe is expanding.




Figure 39.14 Events occur at points P and Q and are observed by an observer at rest in the S frame and another in the S' frame, which is moving to the right with a speed v .

39.5 The Lorentz Transformation Equations

Suppose an event that occurs at some point P is reported by two observers, one at rest in a frame S and another in a frame S' that is moving to the right with speed v as in Figure 39.14. The observer in S reports the event with space–time coordinates (x, y, z, t) , while the observer in S' reports the same event using the coordinates (x', y', z', t') . If two events occur at P and Q, Equation 39.1 predicts that $\Delta x = \Delta x'$, that is, the distance between the two points in space

at which the events occur does not depend on motion of the observer. Because this is contradictory to the notion of length contraction, the Galilean transformation is not valid when v approaches the speed of light. In this section, we state the correct transformation equations that apply for all speeds in the range $0 \leq v < c$.

The equations that are valid for all speeds and enable us to transform coordinates from S to S' are the **Lorentz transformation equations**:

$$x' = \gamma(x - vt) \quad y' = y \quad z' = z \quad t' = \gamma\left(t - \frac{v}{c^2}x\right) \quad (39.11)$$

Lorentz transformation for $S \rightarrow S'$

These transformation equations were developed by Hendrik A. Lorentz (1853–1928) in 1890 in connection with electromagnetism. However, it was Einstein who recognized their physical significance and took the bold step of interpreting them within the framework of the special theory of relativity.

Note the difference between the Galilean and Lorentz time equations. In the Galilean case, $t = t'$, but in the Lorentz case the value for t' assigned to an event by an observer O' in the S' frame in Figure 39.14 depends both on the time t and on the coordinate x as measured by an observer O in the S frame. This is consistent with the notion that an event is characterized by four space–time coordinates (x, y, z, t) . In other words, in relativity, space and time are *not* separate concepts but rather are closely interwoven with each other.

If we wish to transform coordinates in the S' frame to coordinates in the S frame, we simply replace v by $-v$ and interchange the primed and unprimed coordinates in Equations 39.11:

$$x = \gamma(x' + vt') \quad y = y' \quad z = z' \quad t = \gamma\left(t' + \frac{v}{c^2}x'\right) \quad (39.12)$$

Inverse Lorentz transformation for $S' \rightarrow S$

When $v \ll c$, the Lorentz transformation equations should reduce to the Galilean equations. To verify this, note that as v approaches zero, $v/c \ll 1$; thus, $\gamma \rightarrow 1$, and Equations 39.11 reduce to the Galilean space–time transformation equations:

$$x' = x - vt \quad y' = y \quad z' = z \quad t' = t$$

In many situations, we would like to know the difference in coordinates between two events or the time interval between two events as seen by observers O and O' . We can accomplish this by writing the Lorentz equations in a form suitable for describing pairs of events. From Equations 39.11 and 39.12, we can express the differences between the four variables x , x' , t , and t' in the form

$$\left. \begin{aligned} \Delta x' &= \gamma(\Delta x - v \Delta t) \\ \Delta t' &= \gamma\left(\Delta t - \frac{v}{c^2} \Delta x\right) \end{aligned} \right\} S \rightarrow S' \quad (39.13)$$

$$\left. \begin{aligned} \Delta x &= \gamma(\Delta x' + v \Delta t') \\ \Delta t &= \gamma\left(\Delta t' + \frac{v}{c^2} \Delta x'\right) \end{aligned} \right\} S' \rightarrow S \quad (39.14)$$

where $\Delta x' = x'_2 - x'_1$ and $\Delta t' = t'_2 - t'_1$ are the differences measured by observer O' and $\Delta x = x_2 - x_1$ and $\Delta t = t_2 - t_1$ are the differences measured by observer O . (We have not included the expressions for relating the y and z coordinates because they are unaffected by motion along the x direction.⁵)

⁵ Although relative motion of the two frames along the x axis does not change the y and z coordinates of an object, it does change the y and z velocity components of an object moving in either frame, as noted in Section 39.6.

Example 39.6 Simultaneity and Time Dilation Revisited

Use the Lorentz transformation equations in difference form to show that

- (A) simultaneity is not an absolute concept and that
 (B) a moving clock is measured to run more slowly than a clock that is at rest with respect to an observer.

Solution (A) Suppose that two events are simultaneous and separated in space such that $\Delta t' = 0$ and $\Delta x' \neq 0$ according to an observer O' who is moving with speed v relative to O . From the expression for Δt given in Equation 39.14, we see that in this case the time interval Δt measured by observer O is $\Delta t = \gamma v \Delta x' / c^2$. That is, the

time interval for the same two events as measured by O is nonzero, and so the events do not appear to be simultaneous to O .

(B) Suppose that observer O' carries a clock that he uses to measure a time interval $\Delta t'$. He finds that two events occur at the same place in his reference frame ($\Delta x' = 0$) but at different times ($\Delta t' \neq 0$). Observer O' is moving with speed v relative to O , who measures the time interval between the events to be Δt . In this situation, the expression for Δt given in Equation 39.14 becomes $\Delta t = \gamma \Delta t'$. This is the equation for time dilation found earlier (Eq. 39.7), where $\Delta t' = \Delta t_p$ is the proper time measured by the clock carried by observer O' . Thus, O measures the moving clock to run slow.

39.6 The Lorentz Velocity Transformation Equations

Suppose two observers in relative motion with respect to each other are both observing the motion of an object. Previously, we defined an event as occurring at an instant of time. Now, we wish to interpret the “event” as the motion of the object. We know that the Galilean velocity transformation (Eq. 39.2) is valid for low speeds. How do the observers’ measurements of the velocity of the object relate to each other if the speed of the object is close to that of light? Once again S' is our frame moving at a speed v relative to S . Suppose that an object has a velocity component u'_x measured in the S' frame, where

$$u'_x = \frac{dx'}{dt'} \quad (39.15)$$

Using Equation 39.11, we have

$$\begin{aligned} dx' &= \gamma(dx - v dt) \\ dt' &= \gamma\left(dt - \frac{v}{c^2} dx\right) \end{aligned}$$

Substituting these values into Equation 39.15 gives

$$u'_x = \frac{dx'}{dt'} = \frac{dx - v dt}{dt - \frac{v}{c^2} dx} = \frac{\frac{dx}{dt} - v}{1 - \frac{v}{c^2} \frac{dx}{dt}}$$

But dx/dt is just the velocity component u_x of the object measured by an observer in S , and so this expression becomes

$$u'_x = \frac{u_x - v}{1 - \frac{u_x v}{c^2}} \quad (39.16)$$

If the object has velocity components along the y and z axes, the components as measured by an observer in S' are

$$u'_y = \frac{u_y}{\gamma\left(1 - \frac{u_x v}{c^2}\right)} \quad \text{and} \quad u'_z = \frac{u_z}{\gamma\left(1 - \frac{u_x v}{c^2}\right)} \quad (39.17)$$

Lorentz velocity transformation for $S \rightarrow S'$

Note that u'_y and u'_z do not contain the parameter v in the numerator because the relative velocity is along the x axis.

When v is much smaller than c (the nonrelativistic case), the denominator of Equation 39.16 approaches unity, and so $u'_x \approx u_x - v$, which is the Galilean velocity transformation equation. In another extreme, when $u_x = c$, Equation 39.16 becomes

$$u'_x = \frac{c - v}{1 - \frac{cv}{c^2}} = \frac{c \left(1 - \frac{v}{c}\right)}{1 - \frac{v}{c}} = c$$

From this result, we see that a speed measured as c by an observer in S is also measured as c by an observer in S' —independent of the relative motion of S and S' . Note that this conclusion is consistent with Einstein's second postulate—that the speed of light must be c relative to all inertial reference frames. Furthermore, we find that the speed of an object can never be measured as larger than c . That is, the speed of light is the ultimate speed. We return to this point later.

To obtain u_x in terms of u'_x , we replace v by $-v$ in Equation 39.16 and interchange the roles of u_x and u'_x :

$$u_x = \frac{u'_x + v}{1 + \frac{u'_x v}{c^2}} \quad (39.18)$$

▲ PITFALL PREVENTION

39.5 What Can the Observers Agree On?

We have seen several measurements that the two observers O and O' do *not* agree on: (1) the time interval between events that take place in the same position in one of the frames, (2) the distance between two points that remain fixed in one of their frames, (3) the velocity components of a moving particle, and (4) whether two events occurring at different locations in both frames are simultaneous or not. Note that the two observers *can* agree on (1) their relative speed of motion v with respect to each other, (2) the speed c of any ray of light, and (3) the simultaneity of two events which take place at the same position *and* time in some frame.

Quick Quiz 39.8 You are driving on a freeway at a relativistic speed. Straight ahead of you, a technician standing on the ground turns on a searchlight and a beam of light moves exactly vertically upward, as seen by the technician. As you observe the beam of light, you measure the magnitude of the vertical component of its velocity as (a) equal to c (b) greater than c (c) less than c .

Quick Quiz 39.9 Consider the situation in Quick Quiz 39.8 again. If the technician aims the searchlight directly at you instead of upward, you measure the magnitude of the horizontal component of its velocity as (a) equal to c (b) greater than c (c) less than c .

Example 39.7 Relative Velocity of Two Spacecraft

Two spacecraft A and B are moving in opposite directions, as shown in Figure 39.15. An observer on the Earth measures the speed of craft A to be $0.750c$ and the speed of craft B to be $0.850c$. Find the velocity of craft B as observed by the crew on craft A.




Figure 39.15 (Example 39.7) Two spacecraft A and B move in opposite directions. The speed of B relative to A is *less* than c and is obtained from the relativistic velocity transformation equation.

Solution To conceptualize this problem, we carefully identify the observers and the event. The two observers are on the Earth and on spacecraft A. The event is the motion of spacecraft B. Because the problem asks to find an observed velocity, we categorize this problem as one requiring the Lorentz velocity transformation. To analyze the problem, we note that the Earth observer makes two measurements, one of each spacecraft. We identify this observer as being at rest in the S frame. Because the velocity of spacecraft B is what we wish to measure, we identify the speed u_x as $-0.850c$. The velocity of spacecraft A is also the velocity of the observer at rest in the S' frame, which is attached to the spacecraft, relative to the observer at rest in S . Thus, $v = 0.750c$. Now we can obtain the velocity u'_x of craft B relative to craft A by using Equation 39.16:

$$u'_x = \frac{u_x - v}{1 - \frac{u_x v}{c^2}} = \frac{-0.850c - 0.750c}{1 - \frac{(-0.850c)(0.750c)}{c^2}}$$

$$= -0.977c$$

To finalize this problem, note that the negative sign indicates that craft B is moving in the negative x direction as observed by the crew on craft A. Is this consistent with your expectation from Figure 39.15? Note that the speed is less than c . That is, an object whose speed is less than c in one frame of reference must have a speed less than c in any other frame. (If the Galilean velocity transformation equation were used in


this example, we would find that $u'_x = u_x - v = -0.850c - 0.750c = -1.60c$, which is impossible. The Galilean transformation equation does not work in relativistic situations.)

What If? What if the two spacecraft pass each other? Now what is their relative speed?

Answer The calculation using Equation 39.16 involves only the velocities of the two spacecraft and does not depend on their locations. After they pass each other, they have the same velocities, so the velocity of craft B as observed by the crew on craft A is the same, $-0.977c$. The only difference after they pass is that B is receding from A whereas it was approaching A before it passed.

Example 39.8 The Speeding Motorcycle

Imagine a motorcycle moving with a speed $0.80c$ past a stationary observer, as shown in Figure 39.16. If the rider



tosses a ball in the forward direction with a speed of $0.70c$ relative to himself, what is the speed of the ball relative to the stationary observer?

Solution The speed of the motorcycle relative to the stationary observer is $v = 0.80c$. The speed of the ball in the frame of reference of the motorcyclist is $u'_x = 0.70c$. Therefore, the speed u_x of the ball relative to the stationary observer is

$$u_x = \frac{u'_x + v}{1 + \frac{u'_x v}{c^2}} = \frac{0.70c + 0.80c}{1 + \frac{(0.70c)(0.80c)}{c^2}} = 0.96c$$

Figure 39.16 (Example 39.8) A motorcyclist moves past a stationary observer with a speed of $0.80c$ and throws a ball in the direction of motion with a speed of $0.70c$ relative to himself.

Example 39.9 Relativistic Leaders of the Pack

Interactive

Two motorcycle pack leaders named David and Emily are racing at relativistic speeds along perpendicular paths, as shown in Figure 39.17. How fast does Emily recede as seen by David over his right shoulder?

Solution Figure 39.17 represents the situation as seen by a police officer at rest in frame S , who observes the

following:

David: $u_x = 0.75c \quad u_y = 0$
 Emily: $u_x = 0 \quad u_y = -0.90c$

To calculate Emily's speed of recession as seen by David, we take S' to move along with David and then calculate u'_x and




Figure 39.17 (Example 39.9) David moves to the east with a speed $0.75c$ relative to the police officer, and Emily travels south at a speed $0.90c$ relative to the officer.

u'_y for Emily using Equations 39.16 and 39.17:

$$u'_x = \frac{u_x - v}{1 - \frac{u_x v}{c^2}} = \frac{0 - 0.75c}{1 - \frac{(0)(0.75c)}{c^2}} = -0.75c$$

$$u'_y = \frac{u_y}{\gamma \left(1 - \frac{u_x v}{c^2}\right)} = \frac{\sqrt{1 - \frac{(0.75c)^2}{c^2}} (-0.90c)}{\left(1 - \frac{(0)(0.75c)}{c^2}\right)}$$

$$= -0.60c$$

Thus, the speed of Emily as observed by David is

$$u' = \sqrt{(u'_x)^2 + (u'_y)^2} = \sqrt{(-0.75c)^2 + (-0.60c)^2}$$

$$= 0.96c$$

Note that this speed is less than c , as required by the special theory of relativity.



Investigate this situation with various speeds of David and Emily at the Interactive Worked Example link at <http://www.pse6.com>.

39.7 Relativistic Linear Momentum and the Relativistic Form of Newton's Laws

We have seen that in order to describe properly the motion of particles within the framework of the special theory of relativity, we must replace the Galilean transformation equations by the Lorentz transformation equations. Because the laws of physics must remain unchanged under the Lorentz transformation, we must generalize Newton's laws and the definitions of linear momentum and energy to conform to the Lorentz transformation equations and the principle of relativity. These generalized definitions should reduce to the classical (nonrelativistic) definitions for $v \ll c$.

First, recall that the law of conservation of linear momentum states that when two particles (or objects that can be modeled as particles) collide, the total momentum of the isolated system of the two particles remains constant. Suppose that we observe this collision in a reference frame S and confirm that the momentum of the system is conserved. Now imagine that the momenta of the particles are measured by an observer in a second reference frame S' moving with velocity \mathbf{v} relative to the first frame. Using the Lorentz velocity transformation equation and the classical definition of linear momentum, $\mathbf{p} = m\mathbf{u}$ (where \mathbf{u} is the velocity of a particle), we find that linear momentum is *not* measured to be conserved by the observer in S' . However, because the laws of physics are the same in all inertial frames, linear momentum of the system must be conserved in all frames. We have a contradiction. In view of this contradiction and assuming that the Lorentz velocity transformation equation is correct, we must modify the definition of linear momentum to satisfy the following conditions:

- The linear momentum of an isolated system must be conserved in all collisions.
- The relativistic value calculated for the linear momentum \mathbf{p} of a particle must approach the classical value $m\mathbf{u}$ as \mathbf{u} approaches zero.

For any particle, the correct relativistic equation for linear momentum that satisfies these conditions is

$$\mathbf{p} \equiv \frac{m\mathbf{u}}{\sqrt{1 - \frac{u^2}{c^2}}} = \gamma m\mathbf{u} \quad (39.19)$$

where \mathbf{u} is the velocity of the particle and m is the mass of the particle. When u is much less than c , $\gamma = (1 - u^2/c^2)^{-1/2}$ approaches unity and \mathbf{p} approaches $m\mathbf{u}$. Therefore,

PITFALL PREVENTION

39.6 Watch Out for “Relativistic Mass”

Some older treatments of relativity maintained the conservation of momentum principle at high speeds by using a model in which the mass of a particle increases with speed. You might still encounter this notion of “relativistic mass” in your outside reading, especially in older books. Be aware that this notion is no longer widely accepted and mass is considered as *invariant*, independent of speed. The mass of an object in all frames is considered to be the mass as measured by an observer at rest with respect to the object.

Definition of relativistic linear momentum



The speed of light is the speed limit of the Universe. It is the maximum possible speed for energy transfer and for information transfer. Any object with mass must move at a lower speed.

the relativistic equation for \mathbf{p} does indeed reduce to the classical expression when u is much smaller than c .

The relativistic force \mathbf{F} acting on a particle whose linear momentum is \mathbf{p} is defined as

$$\mathbf{F} \equiv \frac{d\mathbf{p}}{dt} \quad (39.20)$$

where \mathbf{p} is given by Equation 39.19. This expression, which is the relativistic form of Newton's second law, is reasonable because it preserves classical mechanics in the limit of low velocities and is consistent with conservation of linear momentum for an isolated system ($\mathbf{F} = 0$) both relativistically and classically.

It is left as an end-of-chapter problem (Problem 69) to show that under relativistic conditions, the acceleration \mathbf{a} of a particle decreases under the action of a constant force, in which case $a \propto (1 - u^2/c^2)^{3/2}$. From this proportionality, we see that as the particle's speed approaches c , the acceleration caused by any finite force approaches zero. Hence, it is impossible to accelerate a particle from rest to a speed $u \geq c$. This argument shows that the speed of light is the ultimate speed, as noted at the end of the preceding section.

Example 39.10 Linear Momentum of an Electron

An electron, which has a mass of 9.11×10^{-31} kg, moves with a speed of $0.750c$. Find its relativistic momentum and compare this value with the momentum calculated from the classical expression.

Solution Using Equation 39.19 with $u = 0.750c$, we have

$$p = \frac{m_e u}{\sqrt{1 - \frac{u^2}{c^2}}}$$

$$p = \frac{(9.11 \times 10^{-31} \text{ kg})(0.750)(3.00 \times 10^8 \text{ m/s})}{\sqrt{1 - \frac{(0.750c)^2}{c^2}}}$$

$$p = 3.10 \times 10^{-22} \text{ kg} \cdot \text{m/s}$$

The classical expression (used incorrectly here) gives

$$p_{\text{classical}} = m_e u = 2.05 \times 10^{-22} \text{ kg} \cdot \text{m/s}$$

Hence, the correct relativistic result is 50% greater than the classical result!

39.8 Relativistic Energy

We have seen that the definition of linear momentum requires generalization to make it compatible with Einstein's postulates. This implies that most likely the definition of kinetic energy must also be modified.

To derive the relativistic form of the work–kinetic energy theorem, let us imagine a particle moving in one dimension along the x axis. A force in the x direction causes the momentum of the particle to change according to Equation 39.20. The work done by the force F on the particle is

$$W = \int_{x_1}^{x_2} F dx = \int_{x_1}^{x_2} \frac{dp}{dt} dx \quad (39.21)$$

In order to perform this integration and find the work done on the particle and the relativistic kinetic energy as a function of u , we first evaluate dp/dt :

$$\frac{dp}{dt} = \frac{d}{dt} \frac{mu}{\sqrt{1 - \frac{u^2}{c^2}}} = \frac{m(du/dt)}{\left(1 - \frac{u^2}{c^2}\right)^{3/2}}$$

Substituting this expression for dp/dt and $dx = u dt$ into Equation 39.21 gives

$$W = \int_0^t \frac{m(du/dt)u dt}{\left(1 - \frac{u^2}{c^2}\right)^{3/2}} = m \int_0^u \frac{u}{\left(1 - \frac{u^2}{c^2}\right)^{3/2}} du$$

where we use the limits 0 and u in the integral because the integration variable has been changed from t to u . We assume that the particle is accelerated from rest to some final speed u . Evaluating the integral, we find that

$$W = \frac{mc^2}{\sqrt{1 - \frac{u^2}{c^2}}} - mc^2 \quad (39.22)$$

Recall from Chapter 7 that the work done by a force acting on a system consisting of a single particle equals the change in kinetic energy of the particle. Because we assumed that the initial speed of the particle is zero, we know that its initial kinetic energy is zero. We therefore conclude that the work W in Equation 39.22 is equivalent to the relativistic kinetic energy K :

$$K = \frac{mc^2}{\sqrt{1 - \frac{u^2}{c^2}}} - mc^2 = \gamma mc^2 - mc^2 = (\gamma - 1)mc^2 \quad (39.23)$$

Relativistic kinetic energy

This equation is routinely confirmed by experiments using high-energy particle accelerators.

At low speeds, where $u/c \ll 1$, Equation 39.23 should reduce to the classical expression $K = \frac{1}{2}mu^2$. We can check this by using the binomial expansion $(1 - \beta^2)^{-1/2} \approx 1 + \frac{1}{2}\beta^2 + \dots$ for $\beta \ll 1$, where the higher-order powers of β are neglected in the expansion. (In treatments of relativity, β is a common symbol used to represent u/c or v/c .) In our case, $\beta = u/c$, so that

$$\gamma = \frac{1}{\sqrt{1 - \frac{u^2}{c^2}}} = \left(1 - \frac{u^2}{c^2}\right)^{-1/2} \approx 1 + \frac{1}{2}\frac{u^2}{c^2}$$

Substituting this into Equation 39.23 gives

$$K \approx \left[\left(1 + \frac{1}{2}\frac{u^2}{c^2}\right) - 1 \right] mc^2 = \frac{1}{2}mu^2 \quad (\text{for } u/c \ll 1)$$

which is the classical expression for kinetic energy. A graph comparing the relativistic and nonrelativistic expressions is given in Figure 39.18. In the relativistic case, the particle speed never exceeds c , regardless of the kinetic energy. The two curves are in good agreement when $u \ll c$.




Figure 39.18 A graph comparing relativistic and nonrelativistic kinetic energy of a moving particle. The energies are plotted as a function of particle speed u . In the relativistic case, u is always less than c .

The constant term mc^2 in Equation 39.23, which is independent of the speed of the particle, is called the **rest energy** E_R of the particle:

Rest energy

$$E_R = mc^2 \quad (39.24)$$

The term γmc^2 , which does depend on the particle speed, is therefore the sum of the kinetic and rest energies. We define γmc^2 to be the **total energy** E :

Total energy = kinetic energy + rest energy

$$E = K + mc^2 \quad (39.25)$$

or

Total energy of a relativistic particle

$$E = \frac{mc^2}{\sqrt{1 - \frac{u^2}{c^2}}} = \gamma mc^2 \quad (39.26)$$

The relationship $E = K + mc^2$ shows that **mass is a form of energy**, where c^2 in the rest energy term is just a constant conversion factor. This expression also shows that a small mass corresponds to an enormous amount of energy, a concept fundamental to nuclear and elementary-particle physics.

In many situations, the linear momentum or energy of a particle is measured rather than its speed. It is therefore useful to have an expression relating the total energy E to the relativistic linear momentum p . This is accomplished by using the expressions $E = \gamma mc^2$ and $p = \gamma mu$. By squaring these equations and subtracting, we can eliminate u (Problem 43). The result, after some algebra, is⁶

Energy–momentum relationship for a relativistic particle

$$E^2 = p^2 c^2 + (mc^2)^2 \quad (39.27)$$

When the particle is at rest, $p = 0$ and so $E = E_R = mc^2$.

In Section 35.1, we introduced the concept of a particle of light, called a **photon**. For particles that have zero mass, such as photons, we set $m = 0$ in Equation 39.27 and find that

$$E = pc \quad (39.28)$$

This equation is an exact expression relating total energy and linear momentum for photons, which always travel at the speed of light (in vacuum).

Finally, note that because the mass m of a particle is independent of its motion, m must have the same value in all reference frames. For this reason, m is often called the **invariant mass**. On the other hand, because the total energy and linear momentum of a particle both depend on velocity, these quantities depend on the reference frame in which they are measured.

When we are dealing with subatomic particles, it is convenient to express their energy in electron volts (Section 25.1) because the particles are usually given this energy by acceleration through a potential difference. The conversion factor, as you recall from Equation 25.5, is

$$1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$$

For example, the mass of an electron is $9.11 \times 10^{-31} \text{ kg}$. Hence, the rest energy of the electron is

$$\begin{aligned} m_e c^2 &= (9.11 \times 10^{-31} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2 = 8.20 \times 10^{-14} \text{ J} \\ &= (8.20 \times 10^{-14} \text{ J})(1 \text{ eV}/1.60 \times 10^{-19} \text{ J}) = 0.511 \text{ MeV} \end{aligned}$$

⁶ One way to remember this relationship is to draw a right triangle having a hypotenuse of length E and legs of lengths pc and mc^2 .

Quick Quiz 39.10 The following *pairs* of energies represent the rest energy and total energy of three different particles: particle 1: E , $2E$; particle 2: E , $3E$; particle 3: $2E$, $4E$. Rank the particles, from greatest to least, according to their (a) mass; (b) kinetic energy; (c) speed.

Example 39.11 The Energy of a Speedy Electron

An electron in a television picture tube typically moves with a speed $u = 0.250c$. Find its total energy and kinetic energy in electron volts.

Solution Using the fact that the rest energy of the electron is 0.511 MeV together with Equation 39.26, we have

$$E = \frac{m_e c^2}{\sqrt{1 - \frac{u^2}{c^2}}} = \frac{0.511 \text{ MeV}}{\sqrt{1 - \frac{(0.250c)^2}{c^2}}}$$

$$= 1.03(0.511 \text{ MeV}) = 0.528 \text{ MeV}$$

This is 3% greater than the rest energy.

We obtain the kinetic energy by subtracting the rest energy from the total energy:

$$K = E - m_e c^2 = 0.528 \text{ MeV} - 0.511 \text{ MeV}$$

$$= 0.017 \text{ MeV}$$

Example 39.12 The Energy of a Speedy Proton

(A) Find the rest energy of a proton in electron volts.

Solution Using Equation 39.24,

$$E_R = m_p c^2 = (1.67 \times 10^{-27} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$$

$$= (1.50 \times 10^{-10} \text{ J}) \left(\frac{1.00 \text{ eV}}{1.60 \times 10^{-19} \text{ J}} \right)$$

$$= 938 \text{ MeV}$$

(B) If the total energy of a proton is three times its rest energy, what is the speed of the proton?

Solution Equation 39.26 gives

$$E = 3m_p c^2 = \frac{m_p c^2}{\sqrt{1 - \frac{u^2}{c^2}}}$$

$$3 = \frac{1}{\sqrt{1 - \frac{u^2}{c^2}}}$$

Solving for u gives

$$\left(1 - \frac{u^2}{c^2} \right) = \frac{1}{9}$$

$$\frac{u^2}{c^2} = \frac{8}{9}$$

$$u = \frac{\sqrt{8}}{3} c = 0.943c = 2.83 \times 10^8 \text{ m/s}$$

(C) Determine the kinetic energy of the proton in electron volts.

Solution From Equation 39.25,

$$K = E - m_p c^2 = 3m_p c^2 - m_p c^2 = 2m_p c^2$$

Because $m_p c^2 = 938 \text{ MeV}$, we see that $K = 1880 \text{ MeV}$.

(D) What is the proton's momentum?

Solution We can use Equation 39.27 to calculate the momentum with $E = 3m_p c^2$:

$$E^2 = p^2 c^2 + (m_p c^2)^2 = (3m_p c^2)^2$$

$$p^2 c^2 = 9(m_p c^2)^2 - (m_p c^2)^2 = 8(m_p c^2)^2$$

$$p = \sqrt{8} \frac{m_p c^2}{c} = \sqrt{8} \frac{(938 \text{ MeV})}{c} = 2650 \text{ MeV}/c$$

The unit of momentum is written MeV/ c , which is a common unit in particle physics.

What If? In classical physics, if the momentum of a particle doubles, the kinetic energy increases by a factor of 4. What happens to the kinetic energy of the speedy proton in this example if its momentum doubles?

Answer Based on what we have seen so far in relativity, it is likely that you would predict that its kinetic energy does not increase by a factor of 4. If the momentum doubles, the new momentum is

$$p_{\text{new}} = 2 \left(\sqrt{8} \frac{m_p c^2}{c} \right) = 4\sqrt{2} \frac{m_p c^2}{c}$$

Using Equation 39.27, we find the square of the new total energy:

$$E_{\text{new}}^2 = p_{\text{new}}^2 c^2 + (m_p c^2)^2$$

$$E_{\text{new}}^2 = \left(4\sqrt{2} \frac{m_p c^2}{c}\right)^2 c^2 + (m_p c^2)^2 = 33(m_p c^2)^2$$

$$E_{\text{new}} = \sqrt{33}(m_p c^2) = 5.7 m_p c^2$$

Now, using Equation 39.25, we find the new kinetic energy:

$$K_{\text{new}} = E_{\text{new}} - m_p c^2 = 5.7 m_p c^2 - m_p c^2 = 4.7 m_p c^2$$

Notice that this is only 2.35 times as large as the kinetic energy we found in part (C), not four times as large. In general, the factor by which the kinetic energy increases if the momentum doubles will depend on the initial momentum, but will approach 4 as the momentum approaches zero. In this latter situation, classical physics correctly describes the situation.

39.9 Mass and Energy

Equation 39.26, $E = \gamma m c^2$, which represents the total energy of a particle, suggests that even when a particle is at rest ($\gamma = 1$) it still possesses enormous energy through its mass. The clearest experimental proof of the equivalence of mass and energy occurs in nuclear and elementary particle interactions in which the conversion of mass into kinetic energy takes place. Because of this, in relativistic situations, we cannot use the principle of conservation of energy as it was outlined in Chapters 7 and 8. We must include rest energy as another form of energy storage.

This concept is important in atomic and nuclear processes, in which the change in mass is a relatively large fraction of the initial mass. For example, in a conventional nuclear reactor, the uranium nucleus undergoes *fission*, a reaction that results in several lighter fragments having considerable kinetic energy. In the case of ^{235}U , which is used as fuel in nuclear power plants, the fragments are two lighter nuclei and a few neutrons. The total mass of the fragments is less than that of the ^{235}U by an amount Δm . The corresponding energy $\Delta m c^2$ associated with this mass difference is exactly equal to the total kinetic energy of the fragments. The kinetic energy is absorbed as the fragments move through water, raising the internal energy of the water. This internal energy is used to produce steam for the generation of electrical power.

Next, consider a basic *fusion* reaction in which two deuterium atoms combine to form one helium atom. The decrease in mass that results from the creation of one helium atom from two deuterium atoms is $\Delta m = 4.25 \times 10^{-29}$ kg. Hence, the corresponding energy that results from one fusion reaction is $\Delta m c^2 = 3.83 \times 10^{-12}$ J = 23.9 MeV. To appreciate the magnitude of this result, if only 1 g of deuterium is converted to helium, the energy released is on the order of 10^{12} J! At the year 2003 cost of electrical energy, this would be worth about \$30 000. We shall present more details of these nuclear processes in Chapter 45 of the extended version of this textbook.

Example 39.13 Mass Change in a Radioactive Decay

The ^{216}Po nucleus is unstable and exhibits radioactivity (Chapter 44). It decays to ^{212}Pb by emitting an alpha particle, which is a helium nucleus, ^4He . Find

- (A) the mass change in this decay and
- (B) the energy that this represents.

Solution Using values in Table A.3, we see that the initial and final masses are

$$m_i = m(^{216}\text{Po}) = 216.001\,905\text{ u}$$

$$m_f = m(^{212}\text{Pb}) + m(^4\text{He}) = 211.991\,888\text{ u} + 4.002\,603\text{ u} \\ = 215.994\,491\text{ u}$$

Thus, the mass change is

$$\Delta m = 216.001\,905\text{ u} - 215.994\,491\text{ u} = 0.007\,414\text{ u} \\ = 1.23 \times 10^{-29}\text{ kg}$$

(B) The energy associated with this mass change is

$$E = \Delta m c^2 = (1.23 \times 10^{-29}\text{ kg})(3.00 \times 10^8\text{ m/s})^2 \\ = 1.11 \times 10^{-12}\text{ J} = 6.92\text{ MeV}$$

This energy appears as the kinetic energy of the alpha particle and the ^{212}Pb nucleus after the decay.

39.10 The General Theory of Relativity

Up to this point, we have sidestepped a curious puzzle. Mass has two seemingly different properties: a *gravitational attraction* for other masses and an *inertial* property that represents a resistance to acceleration. To designate these two attributes, we use the subscripts g and i and write

$$\begin{aligned} \text{Gravitational property} \quad F_g &= m_g g \\ \text{Inertial property} \quad \sum F &= m_i a \end{aligned}$$

The value for the gravitational constant G was chosen to make the magnitudes of m_g and m_i numerically equal. Regardless of how G is chosen, however, the strict proportionality of m_g and m_i has been established experimentally to an extremely high degree: a few parts in 10^{12} . Thus, it appears that gravitational mass and inertial mass may indeed be exactly proportional.

But why? They seem to involve two entirely different concepts: a force of mutual gravitational attraction between two masses, and the resistance of a single mass to being accelerated. This question, which puzzled Newton and many other physicists over the years, was answered by Einstein in 1916 when he published his theory of gravitation, known as the *general theory of relativity*. Because it is a mathematically complex theory, we offer merely a hint of its elegance and insight.

In Einstein's view, the dual behavior of mass was evidence for a very intimate and basic connection between the two behaviors. He pointed out that no mechanical experiment (such as dropping an object) could distinguish between the two situations illustrated in Figures 39.19a and 39.19b. In Figure 39.19a, a person is standing in an elevator on the surface of a planet, and feels pressed into the floor, due to the gravitational force. In Figure 39.19b, the person is in an elevator in empty space accelerating upward with $a = g$. The person feels pressed into the floor with the same force as in Figure 39.19a. In each case, an object released by the observer undergoes a downward acceleration of magnitude g relative to the floor. In Figure 39.19a, the person is in an inertial frame in a gravitational field. In Figure 39.19b, the person is in a noninertial frame accelerating in gravity-free space. Einstein's claim is that these two situations are completely equivalent.




Figure 39.19 (a) The observer is at rest in a uniform gravitational field \mathbf{g} , directed downward. (b) The observer is in a region where gravity is negligible, but the frame is accelerated by an external force \mathbf{F} that produces an acceleration \mathbf{g} directed upward. According to Einstein, the frames of reference in parts (a) and (b) are equivalent in every way. No local experiment can distinguish any difference between the two frames. (c) In the accelerating frame, a ray of light would appear to bend downward due to the acceleration of the elevator. (d) If parts (a) and (b) are truly equivalent, as Einstein proposed, then part (c) suggests that a ray of light would bend downward in a gravitational field.

Einstein carried this idea further and proposed that *no* experiment, mechanical or otherwise, could distinguish between the two cases. This extension to include all phenomena (not just mechanical ones) has interesting consequences. For example, suppose that a light pulse is sent horizontally across an elevator that is accelerating upward in empty space, as in Figure 39.19c. From the point of view of an observer in an inertial frame outside of the elevator, the light travels in a straight line while the floor of the elevator accelerates upward. According to the observer on the elevator, however, the trajectory of the light pulse bends downward as the floor of the elevator (and the observer) accelerates upward. Therefore, based on the equality of parts (a) and (b) of the figure for all phenomena, Einstein proposed that **a beam of light should also be bent downward by a gravitational field**, as in Figure 39.19d. Experiments have verified the effect, although the bending is small. A laser aimed at the horizon falls less than 1 cm after traveling 6 000 km. (No such bending is predicted in Newton's theory of gravitation.)

The two postulates of Einstein's **general theory of relativity** are

Postulates of the general theory of relativity


- All the laws of nature have the same form for observers in any frame of reference, whether accelerated or not.
- In the vicinity of any point, a gravitational field is equivalent to an accelerated frame of reference in the absence of gravitational effects. (This is the **principle of equivalence**.)

One interesting effect predicted by the general theory is that time is altered by gravity. A clock in the presence of gravity runs slower than one located where gravity is negligible. Consequently, the frequencies of radiation emitted by atoms in the presence of a strong gravitational field are *red-shifted* to lower frequencies when compared with the same emissions in the presence of a weak field. This gravitational red shift has been detected in spectral lines emitted by atoms in massive stars. It has also been verified on the Earth by comparing the frequencies of gamma rays emitted from nuclei separated vertically by about 20 m.

The second postulate suggests that a gravitational field may be “transformed away” at any point if we choose an appropriate accelerated frame of reference—a freely falling one. Einstein developed an ingenious method of describing the acceleration necessary to make the gravitational field “disappear.” He specified a concept, the *curvature of space–time*, that describes the gravitational effect at every point. In fact, the curvature of space–time completely replaces Newton's gravitational theory. According to Einstein, there is no such thing as a gravitational force. Rather, the presence of a mass causes a curvature of space–time in the vicinity of the mass, and this curvature dictates the space–time path that all freely moving objects must follow. In 1979, John Wheeler summarized Einstein's general theory of relativity in a single sentence: “Space tells matter how to move and matter tells space how to curve.”

As an example of the effects of curved space–time, imagine two travelers moving on parallel paths a few meters apart on the surface of the Earth and maintaining an exact northward heading along two longitude lines. As they observe each other near the equator, they will claim that their paths are exactly parallel. As they approach the North Pole, however, they notice that they are moving closer together, and they will actually meet at the North Pole. Thus, they will claim that they moved along parallel paths, but moved toward each other, *as if there were an attractive force between them*. They will make this conclusion based on their everyday experience of moving on flat surfaces. From our mental representation, however, we realize that they are walking on a curved surface, and it is the geometry of the curved surface that causes them to converge, rather than an attractive force. In a similar way, general relativity replaces the notion of forces with the movement of objects through curved space–time.

One prediction of the general theory of relativity is that a light ray passing near the Sun should be deflected in the curved space–time created by the Sun's mass. This prediction was confirmed when astronomers detected the bending of starlight near the



Courtesy of NASA

Einstein's cross. The four bright spots are images of the same galaxy that have been bent around a massive object located between the galaxy and the Earth. The massive object acts like a lens, causing the rays of light that were diverging from the distant galaxy to converge on the Earth. (If the intervening massive object had a uniform mass distribution, we would see a bright ring instead of four spots.)




Figure 39.20 Deflection of starlight passing near the Sun. Because of this effect, the Sun or some other remote object can act as a *gravitational lens*. In his general theory of relativity, Einstein calculated that starlight just grazing the Sun’s surface should be deflected by an angle of 1.75 s of arc.

Sun during a total solar eclipse that occurred shortly after World War I (Fig. 39.20). When this discovery was announced, Einstein became an international celebrity.

If the concentration of mass becomes very great, as is believed to occur when a large star exhausts its nuclear fuel and collapses to a very small volume, a **black hole** may form. Here, the curvature of space–time is so extreme that, within a certain distance from the center of the black hole, all matter and light become trapped, as discussed in Section 13.7.

SUMMARY

The two basic postulates of the special theory of relativity are

- The laws of physics must be the same in all inertial reference frames.
- The speed of light in vacuum has the same value, $c = 3.00 \times 10^8$ m/s, in all inertial frames, regardless of the velocity of the observer or the velocity of the source emitting the light.

Three consequences of the special theory of relativity are

- Events that are measured to be simultaneous for one observer are not necessarily measured to be simultaneous for another observer who is in motion relative to the first.
- Clocks in motion relative to an observer are measured to run slower by a factor $\gamma = (1 - v^2/c^2)^{-1/2}$. This phenomenon is known as **time dilation**.
- The length of objects in motion are measured to be contracted in the direction of motion by a factor $1/\gamma = (1 - v^2/c^2)^{1/2}$. This phenomenon is known as **length contraction**.

To satisfy the postulates of special relativity, the Galilean transformation equations must be replaced by the **Lorentz transformation equations**:

$$x' = \gamma(x - vt) \quad y' = y \quad z' = z \quad t' = \gamma\left(t - \frac{v}{c^2}x\right) \quad (39.11)$$

where $\gamma = (1 - v^2/c^2)^{-1/2}$ and the S' frame moves in the x direction relative to the S frame.

The relativistic form of the **velocity transformation equation** is

$$u'_x = \frac{u_x - v}{1 - \frac{u_x v}{c^2}} \quad (39.16)$$

where u_x is the speed of an object as measured in the S frame and u'_x is its speed measured in the S' frame.

 Take a practice test for this chapter by clicking on the Practice Test link at <http://www.pse6.com>.

The relativistic expression for the **linear momentum** of a particle moving with a velocity \mathbf{u} is

$$\mathbf{p} \equiv \frac{m\mathbf{u}}{\sqrt{1 - \frac{u^2}{c^2}}} = \gamma m\mathbf{u} \quad (39.19)$$

The relativistic expression for the **kinetic energy** of a particle is

$$K = \frac{mc^2}{\sqrt{1 - \frac{u^2}{c^2}}} - mc^2 = (\gamma - 1)mc^2 \quad (39.23)$$

The constant term mc^2 in Equation 39.23 is called the **rest energy** E_R of the particle:

$$E_R = mc^2 \quad (39.24)$$

The total energy E of a particle is given by

$$E = \frac{mc^2}{\sqrt{1 - \frac{u^2}{c^2}}} = \gamma mc^2 \quad (39.26)$$

The relativistic linear momentum of a particle is related to its total energy through the equation

$$E^2 = p^2c^2 + (mc^2)^2 \quad (39.27)$$

QUESTIONS

- What two speed measurements do two observers in relative motion always agree on?
- A spacecraft with the shape of a sphere moves past an observer on Earth with a speed $0.5c$. What shape does the observer measure for the spacecraft as it moves past?
- The speed of light in water is 230 Mm/s . Suppose an electron is moving through water at 250 Mm/s . Does this violate the principle of relativity?
- Two identical clocks are synchronized. One is then put in orbit directed eastward around the Earth while the other remains on the Earth. Which clock runs slower? When the moving clock returns to the Earth, are the two still synchronized?
- Explain why it is necessary, when defining the length of a rod, to specify that the positions of the ends of the rod are to be measured simultaneously.
- A train is approaching you at very high speed as you stand next to the tracks. Just as an observer on the train passes you, you both begin to play the same Beethoven symphony on portable compact disc players. (a) According to you, whose CD player finishes the symphony first? (b) **What If?** According to the observer on the train, whose CD player finishes the symphony first? (c) Whose CD player really finishes the symphony first?
- List some ways our day-to-day lives would change if the speed of light were only 50 m/s .
- Does saying that a moving clock runs slower than a stationary one imply that something is physically unusual about the moving clock?
- How is acceleration indicated on a space–time graph?
- A particle is moving at a speed less than $c/2$. If the speed of the particle is doubled, what happens to its momentum?
- Give a physical argument that shows that it is impossible to accelerate an object of mass m to the speed of light, even with a continuous force acting on it.
- The upper limit of the speed of an electron is the speed of light c . Does that mean that the momentum of the electron has an upper limit?
- Because mass is a measure of energy, can we conclude that the mass of a compressed spring is greater than the mass of the same spring when it is not compressed?
- It is said that Einstein, in his teenage years, asked the question, “What would I see in a mirror if I carried it in my hands and ran at the speed of light?” How would you answer this question?
- Some distant astronomical objects, called quasars, are receding from us at half the speed of light (or greater). What is the speed of the light we receive from these quasars?
- Photons of light have zero mass. How is it possible that they have momentum?
- “Newtonian mechanics correctly describes objects moving at ordinary speeds and relativistic mechanics correctly describes objects moving very fast.” “Relativistic mechanics must make a smooth transition as it reduces to Newtonian mechanics in a case where the speed of an object becomes small compared to the speed of light.” Argue for or against each of these two statements.




Figure Q39.18

18. Two cards have straight edges. Suppose that the top edge of one card crosses the bottom edge of another card at a small angle, as in Figure Q39.18a. A person slides the cards together at a moderately high speed. In what direction does the intersection point of the edges move? Show that it can move at a speed greater than the speed of light.
- A small flashlight is suspended in a horizontal plane and set into rapid rotation. Show that the spot of light it produces on a distant screen can move across the screen at a speed greater than the speed of light. (If you use a laser pointer, as in Figure Q39.18b, make sure the direct laser light cannot enter a person's eyes.) Argue that these experiments do not invalidate the principle that no material, no energy, and no information can move faster than light moves in a vacuum.
19. Describe how the results of Example 39.7 would change if, instead of fast space vehicles, two ordinary cars were approaching each other at highway speeds.
20. Two objects are identical except that one is hotter than the other. Compare how they respond to identical forces.
21. With regard to reference frames, how does general relativity differ from special relativity?
22. Two identical clocks are in the same house, one upstairs in a bedroom, and the other downstairs in the kitchen. Which clock runs more slowly? Explain.
23. *A thought experiment.* Imagine ants living on a merry-go-round turning at relativistic speed, which is their two-dimensional world. From measurements on small circles they are thoroughly familiar with the number π . When they measure the circumference of their world, and divide it by the diameter, they expect to calculate the number $\pi = 3.14159. . .$. We see the merry-go-round turning at relativistic speed. From our point of view, the ants' measuring rods on the circumference are experiencing length contraction in the tangential direction; hence the ants will need some extra rods to fill that entire distance. The rods measuring the diameter, however, do not contract, because their motion is perpendicular to their lengths. As a result, the computed ratio does not agree with the number π . If you were an ant, you would say that the rest of the universe is spinning in circles, and your disk is stationary. What possible explanation can you then give for the discrepancy, in light of the general theory of relativity?

PROBLEMS

- 1, 2, 3 = straightforward, intermediate, challenging = full solution available in the *Student Solutions Manual and Study Guide*
-  = coached solution with hints available at <http://www.pse6.com>  = computer useful in solving problem
-  = paired numerical and symbolic problems

Section 39.1 The Principle of Galilean Relativity

- A 2000-kg car moving at 20.0 m/s collides and locks together with a 1500-kg car at rest at a stop sign. Show that momentum is conserved in a reference frame moving at 10.0 m/s in the direction of the moving car.
- A ball is thrown at 20.0 m/s inside a boxcar moving along the tracks at 40.0 m/s. What is the speed of the ball

relative to the ground if the ball is thrown (a) forward (b) backward (c) out the side door?

- In a laboratory frame of reference, an observer notes that Newton's second law is valid. Show that it is also valid for an observer moving at a constant speed, small compared with the speed of light, relative to the laboratory frame.

4. Show that Newton's second law is *not* valid in a reference frame moving past the laboratory frame of Problem 3 with a constant acceleration.

Section 39.2 The Michelson–Morley Experiment

Section 39.3 Einstein's Principle of Relativity

Section 39.4 Consequences of the Special Theory of Relativity

Problem 43 in Chapter 4 can be assigned with this section.

5. How fast must a meter stick be moving if its length is measured to shrink to 0.500 m?
6. At what speed does a clock move if it is measured to run at a rate that is half the rate of a clock at rest with respect to an observer?
7. An astronaut is traveling in a space vehicle that has a speed of $0.500c$ relative to the Earth. The astronaut measures her pulse rate at 75.0 beats per minute. Signals generated by the astronaut's pulse are radioed to Earth when the vehicle is moving in a direction perpendicular to the line that connects the vehicle with an observer on the Earth. (a) What pulse rate does the Earth observer measure? (b) **What If?** What would be the pulse rate if the speed of the space vehicle were increased to $0.990c$?
8. An astronomer on Earth observes a meteoroid in the southern sky approaching the Earth at a speed of $0.800c$. At the time of its discovery the meteoroid is 20.0 ly from the Earth. Calculate (a) the time interval required for the meteoroid to reach the Earth as measured by the Earth-bound astronomer, (b) this time interval as measured by a tourist on the meteoroid, and (c) the distance to the Earth as measured by the tourist.
9. An atomic clock moves at 1 000 km/h for 1.00 h as measured by an identical clock on the Earth. How many nanoseconds slow will the moving clock be compared with the Earth clock, at the end of the 1.00-h interval?
10. A muon formed high in the Earth's atmosphere travels at speed $v = 0.990c$ for a distance of 4.60 km before it decays into an electron, a neutrino, and an antineutrino ($\mu^- \rightarrow e^- + \nu + \bar{\nu}$). (a) How long does the muon live, as measured in its reference frame? (b) How far does the Earth travel, as measured in the frame of the muon?
11.  A spacecraft with a proper length of 300 m takes $0.750 \mu\text{s}$ to pass an Earth observer. Determine the speed of the spacecraft as measured by the Earth observer.
12. (a) An object of proper length L_p takes a time interval Δt to pass an Earth observer. Determine the speed of the object as measured by the Earth observer. (b) A column of tanks, 300 m long, takes 75.0 s to pass a child waiting at a street corner on her way to school. Determine the speed of the armored vehicles. (c) Show that the answer to part (a) includes the answer to Problem 11 as a special case, and includes the answer to part (b) as another special case.
13. **Review problem.** In 1963 Mercury astronaut Gordon Cooper orbited the Earth 22 times. The press stated that for each orbit he aged 2 millionths of a second less than he would have if he had remained on the Earth. (a) Assuming that he was 160 km above the Earth in a circular orbit, determine the time difference between someone on the Earth and the orbiting astronaut for the 22 orbits. You will need to use the approximation $\sqrt{1-x} \approx 1-x/2$, for small x . (b) Did the press report accurate information? Explain.
14. For what value of v does $\gamma = 1.0100$? Observe that for speeds lower than this value, time dilation and length contraction are effects amounting to less than 1%.
15. A friend passes by you in a spacecraft traveling at a high speed. He tells you that his craft is 20.0 m long and that the identically constructed craft you are sitting in is 19.0 m long. According to your observations, (a) how long is your spacecraft, (b) how long is your friend's craft, and (c) what is the speed of your friend's craft?
16. The identical twins Speedo and Goslo join a migration from the Earth to Planet X. It is 20.0 ly away in a reference frame in which both planets are at rest. The twins, of the same age, depart at the same time on different spacecraft. Speedo's craft travels steadily at $0.950c$, and Goslo's at $0.750c$. Calculate the age difference between the twins after Goslo's spacecraft lands on Planet X. Which twin is the older?
17. An interstellar space probe is launched from the Earth. After a brief period of acceleration it moves with a constant velocity, with a magnitude of 70.0% of the speed of light. Its nuclear-powered batteries supply the energy to keep its data transmitter active continuously. The batteries have a lifetime of 15.0 yr as measured in a rest frame. (a) How long do the batteries on the space probe last as measured by Mission Control on the Earth? (b) How far is the probe from the Earth when its batteries fail, as measured by Mission Control? (c) How far is the probe from the Earth when its batteries fail, as measured by its built-in trip odometer? (d) For what total time interval after launch are data received from the probe by Mission Control? Note that radio waves travel at the speed of light and fill the space between the probe and the Earth at the time of battery failure.
18. **Review problem.** An alien civilization occupies a brown dwarf, nearly stationary relative to the Sun, several lightyears away. The extraterrestrials have come to love original broadcasts of *I Love Lucy*, on our television channel 2, at carrier frequency 57.0 MHz. Their line of sight to us is in the plane of the Earth's orbit. Find the difference between the highest and lowest frequencies they receive due to the Earth's orbital motion around the Sun.
19. Police radar detects the speed of a car (Fig. P39.19) as follows. Microwaves of a precisely known frequency are broadcast toward the car. The moving car reflects the microwaves with a Doppler shift. The reflected waves are received and combined with an attenuated version of the transmitted wave. Beats occur between the two microwave signals. The beat frequency is measured. (a) For an electromagnetic wave reflected back to its source from a mirror approaching at speed v , show that the reflected

wave has frequency

$$f = f_{\text{source}} \frac{c + v}{c - v}$$

where f_{source} is the source frequency. (b) When v is much less than c , the beat frequency is much smaller than the transmitted frequency. In this case use the approximation $f + f_{\text{source}} \approx 2f_{\text{source}}$ and show that the beat frequency can be written as $f_{\text{beat}} = 2v/\lambda$. (c) What beat frequency is measured for a car speed of 30.0 m/s if the microwaves have frequency 10.0 GHz? (d) If the beat frequency measurement is accurate to ± 5 Hz, how accurate is the velocity measurement?




Figure P39.19

20. *The red shift.* A light source recedes from an observer with a speed v_{source} that is small compared with c . (a) Show that the fractional shift in the measured wavelength is given by the approximate expression

$$\frac{\Delta\lambda}{\lambda} \approx \frac{v_{\text{source}}}{c}$$

This phenomenon is known as the red shift, because the visible light is shifted toward the red. (b) Spectroscopic measurements of light at $\lambda = 397$ nm coming from a galaxy in Ursa Major reveal a red shift of 20.0 nm. What is the recessional speed of the galaxy?

21. A physicist drives through a stop light. When he is pulled over, he tells the police officer that the Doppler shift made the red light of wavelength 650 nm appear green to him, with a wavelength of 520 nm. The police officer writes out a traffic citation for speeding. How fast was the physicist traveling, according to his own testimony?

Section 39.5 The Lorentz Transformation Equations

22. Suzanne observes two light pulses to be emitted from the same location, but separated in time by 3.00 μs . Mark sees

the emission of the same two pulses separated in time by 9.00 μs . (a) How fast is Mark moving relative to Suzanne? (b) According to Mark, what is the separation in space of the two pulses?

23. A moving rod is observed to have a length of 2.00 m and to be oriented at an angle of 30.0° with respect to the direction of motion, as shown in Figure P39.23. The rod has a speed of $0.995c$. (a) What is the proper length of the rod? (b) What is the orientation angle in the proper frame?




Figure P39.23

24. An observer in reference frame S sees two events as simultaneous. Event A occurs at the point (50.0 m, 0, 0) at the instant 9:00:00 Universal time, 15 January 2004. Event B occurs at the point (150 m, 0, 0) at the same moment. A second observer, moving past with a velocity of $0.800c\hat{i}$, also observes the two events. In her reference frame S' , which event occurred first and what time interval elapsed between the events?
25. A red light flashes at position $x_R = 3.00$ m and time $t_R = 1.00 \times 10^{-9}$ s, and a blue light flashes at $x_B = 5.00$ m and $t_B = 9.00 \times 10^{-9}$ s, all measured in the S reference frame. Reference frame S' has its origin at the same point as S at $t = t' = 0$; frame S' moves uniformly to the right. Both flashes are observed to occur at the same place in S' . (a) Find the relative speed between S and S' . (b) Find the location of the two flashes in frame S' . (c) At what time does the red flash occur in the S' frame?

Section 39.6 The Lorentz Velocity Transformation Equations

26. A Klingon spacecraft moves away from the Earth at a speed of $0.800c$ (Fig. P39.26). The starship *Enterprise* pursues at a speed of $0.900c$ relative to the Earth. Observers on the Earth see the *Enterprise* overtaking the Klingon craft at a relative speed of $0.100c$. With what speed is the *Enterprise* overtaking the Klingon craft as seen by the crew of the *Enterprise*?




Figure P39.26

27. Two jets of material from the center of a radio galaxy are ejected in opposite directions. Both jets move at $0.750c$

relative to the galaxy. Determine the speed of one jet relative to the other.

28. A spacecraft is launched from the surface of the Earth with a velocity of $0.600c$ at an angle of 50.0° above the horizontal positive x axis. Another spacecraft is moving past, with a velocity of $0.700c$ in the negative x direction. Determine the magnitude and direction of the velocity of the first spacecraft as measured by the pilot of the second spacecraft.

Section 39.7 Relativistic Linear Momentum and the Relativistic Form of Newton's Laws

29. Calculate the momentum of an electron moving with a speed of (a) $0.0100c$, (b) $0.500c$, and (c) $0.900c$.
30. The nonrelativistic expression for the momentum of a particle, $p = mu$, agrees with experiment if $u \ll c$. For what speed does the use of this equation give an error in the momentum of (a) 1.00% and (b) 10.0%?
31. A golf ball travels with a speed of 90.0 m/s. By what fraction does its relativistic momentum magnitude p differ from its classical value mu ? That is, find the ratio $(p - mu)/mu$.
32. Show that the speed of an object having momentum of magnitude p and mass m is

$$u = \frac{c}{\sqrt{1 + (mc/p)^2}}$$

33.  An unstable particle at rest breaks into two fragments of unequal mass. The mass of the first fragment is 2.50×10^{-28} kg, and that of the other is 1.67×10^{-27} kg. If the lighter fragment has a speed of $0.893c$ after the breakup, what is the speed of the heavier fragment?

Section 39.8 Relativistic Energy

34. Determine the energy required to accelerate an electron from (a) $0.500c$ to $0.900c$ and (b) $0.900c$ to $0.990c$.
35. A proton in a high-energy accelerator moves with a speed of $c/2$. Use the work–kinetic energy theorem to find the work required to increase its speed to (a) $0.750c$ and (b) $0.995c$.
36. Show that, for any object moving at less than one-tenth the speed of light, the relativistic kinetic energy agrees with the result of the classical equation $K = \frac{1}{2}mu^2$ to within less than 1%. Thus for most purposes, the classical equation is good enough to describe these objects, whose motion we call *nonrelativistic*.
37. Find the momentum of a proton in MeV/c units assuming its total energy is twice its rest energy.
38. Find the kinetic energy of a 78.0 -kg spacecraft launched out of the solar system with speed 106 km/s by using (a) the classical equation $K = \frac{1}{2}mu^2$. (b) **What If?** Calculate its kinetic energy using the relativistic equation.
39.  A proton moves at $0.950c$. Calculate its (a) rest energy, (b) total energy, and (c) kinetic energy.
40. A cube of steel has a volume of 1.00 cm³ and a mass of 8.00 g when at rest on the Earth. If this cube is now given a speed $u = 0.900c$, what is its density as measured by a

stationary observer? Note that relativistic density is defined as E_R/c^2V .

41. An unstable particle with a mass of 3.34×10^{-27} kg is initially at rest. The particle decays into two fragments that fly off along the x axis with velocity components $0.987c$ and $-0.868c$. Find the masses of the fragments. (*Suggestion:* Conserve both energy and momentum.)
42. An object having mass 900 kg and traveling at speed $0.850c$ collides with a stationary object having mass 1400 kg. The two objects stick together. Find (a) the speed and (b) the mass of the composite object.
43. Show that the energy–momentum relationship $E^2 = p^2c^2 + (mc^2)^2$ follows from the expressions $E = \gamma mc^2$ and $p = \gamma mu$.
44. In a typical color television picture tube, the electrons are accelerated through a potential difference of 25000 V. (a) What speed do the electrons have when they strike the screen? (b) What is their kinetic energy in joules?
45. Consider electrons accelerated to an energy of 20.0 GeV in the 3.00 -km-long Stanford Linear Accelerator. (a) What is the γ factor for the electrons? (b) What is their speed? (c) How long does the accelerator appear to them?
46. Compact high-power lasers can produce a 2.00 -J light pulse of duration 100 fs, focused to a spot 1 μm in diameter. (See Mourou and Umstadter, “Extreme Light,” *Scientific American*, May 2002, page 81.) The electric field in the light accelerates electrons in the target material to near the speed of light. (a) What is the average power of the laser during the pulse? (b) How many electrons can be accelerated to $0.9999c$ if 0.0100% of the pulse energy is converted into energy of electron motion?
47. A pion at rest ($m_\pi = 273m_e$) decays to a muon ($m_\mu = 207m_e$) and an antineutrino ($m_{\bar{\nu}} \approx 0$). The reaction is written $\pi^- \rightarrow \mu^- + \bar{\nu}$. Find the kinetic energy of the muon and the energy of the antineutrino in electron volts. (*Suggestion:* Conserve both energy and momentum.)
48. According to observer A, two objects of equal mass and moving along the x axis collide head on and stick to each other. Before the collision, this observer measures that object 1 moves to the right with a speed of $3c/4$, while object 2 moves to the left with the same speed. According to observer B, however, object 1 is initially at rest. (a) Determine the speed of object 2 as seen by observer B. (b) Compare the total initial energy of the system in the two frames of reference.

Section 39.9 Mass and Energy

49. Make an order-of-magnitude estimate of the ratio of mass increase to the original mass of a flag, as you run it up a flagpole. In your solution explain what quantities you take as data and the values you estimate or measure for them.
50. When 1.00 g of hydrogen combines with 8.00 g of oxygen, 9.00 g of water is formed. During this chemical reaction, 2.86×10^5 J of energy is released. How much mass do the constituents of this reaction lose? Is the loss of mass likely to be detectable?
51. In a nuclear power plant the fuel rods last 3 yr before they are replaced. If a plant with rated thermal power 1.00 GW

operates at 80.0% capacity for 3.00 yr, what is the loss of mass of the fuel?

52. **Review problem.** The total volume of water in the oceans is approximately $1.40 \times 10^9 \text{ km}^3$. The density of sea water is 1030 kg/m^3 , and the specific heat of the water is $4186 \text{ J/(kg} \cdot \text{ }^\circ\text{C)}$. Find the increase in mass of the oceans produced by an increase in temperature of 10.0°C .
53. The power output of the Sun is $3.77 \times 10^{26} \text{ W}$. How much mass is converted to energy in the Sun each second?
54. A gamma ray (a high-energy photon) can produce an electron (e^-) and a positron (e^+) when it enters the electric field of a heavy nucleus: $\gamma \rightarrow e^+ + e^-$. What minimum gamma-ray energy is required to accomplish this task? (Note: The masses of the electron and the positron are equal.)

Section 39.10 The General Theory of Relativity

55. An Earth satellite used in the global positioning system moves in a circular orbit with period 11 h 58 min. (a) Determine the radius of its orbit. (b) Determine its speed. (c) The satellite contains an oscillator producing the principal nonmilitary GPS signal. Its frequency is 1575.42 MHz in the reference frame of the satellite. When it is received on the Earth’s surface, what is the fractional change in this frequency due to time dilation, as described by special relativity? (d) The gravitational blue shift of the frequency according to general relativity is a separate effect. The magnitude of that fractional change is given by

$$\frac{\Delta f}{f} = \frac{\Delta U_g}{mc^2}$$

where ΔU_g is the change in gravitational potential energy of an object–Earth system when the object of mass m is moved between the two points at which the signal is observed. Calculate this fractional change in frequency. (e) What is the overall fractional change in frequency? Superposed on both of these relativistic effects is a Doppler shift that is generally much larger. It can be a red shift or a blue shift, depending on the motion of a particular satellite relative to a GPS receiver (Fig. P39.55).



Figure P39.55 This global positioning system (GPS) receiver incorporates relativistically corrected time calculations in its analysis of signals it receives from orbiting satellites. This allows the unit to determine its position on the Earth’s surface to within a few meters. If these corrections were not made, the location error would be about 1 km.

Additional Problems

56. An astronaut wishes to visit the Andromeda galaxy, making a one-way trip that will take 30.0 yr in the spacecraft’s frame of reference. Assume that the galaxy is $2.00 \times 10^6 \text{ ly}$ away and that the astronaut’s speed is constant. (a) How fast must he travel relative to the Earth? (b) What will be the kinetic energy of his 1000-metric-ton spacecraft? (c) What is the cost of this energy if it is purchased at a typical consumer price for electric energy: \$0.130/kWh?
57. The cosmic rays of highest energy are protons that have kinetic energy on the order of 10^{13} MeV . (a) How long would it take a proton of this energy to travel across the Milky Way galaxy, having a diameter $\sim 10^5 \text{ ly}$, as measured in the proton’s frame? (b) From the point of view of the proton, how many kilometers across is the galaxy?
58. An electron has a speed of $0.750c$. (a) Find the speed of a proton that has the same kinetic energy as the electron. (b) **What If?** Find the speed of a proton that has the same momentum as the electron.
59. Ted and Mary are playing a game of catch in frame S' , which is moving at $0.600c$ with respect to frame S , while Jim, at rest in frame S , watches the action (Fig. P39.59). Ted throws the ball to Mary at $0.800c$ (according to Ted) and their separation (measured in S') is $1.80 \times 10^{12} \text{ m}$. (a) According to Mary, how fast is the ball moving? (b) According to Mary, how long does it take the ball to reach her? (c) According to Jim, how far apart are Ted and Mary, and how fast is the ball moving? (d) According to Jim, how long does it take the ball to reach Mary?




Figure P39.59

60. A rechargeable AA battery with a mass of 25.0 g can supply a power of 1.20 W for 50.0 min. (a) What is the difference in mass between a charged and an uncharged battery? (b) What fraction of the total mass is this mass difference?
61. The net nuclear fusion reaction inside the Sun can be written as $4 \text{ }^1\text{H} \rightarrow \text{}^4\text{He} + \Delta E$. The rest energy of each hydrogen atom is 938.78 MeV and the rest energy of the helium-4 atom is 3728.4 MeV. Calculate the percentage of the starting mass that is transformed to other forms of energy.
62. An object disintegrates into two fragments. One of the fragments has mass $1.00 \text{ MeV}/c^2$ and momentum $1.75 \text{ MeV}/c$ in the positive x direction. The other fragment has mass $1.50 \text{ MeV}/c^2$ and momentum $2.00 \text{ MeV}/c$ in the positive y direction. Find (a) the mass and (b) the speed of the original object.

63. An alien spaceship traveling at $0.600c$ toward the Earth launches a landing craft with an advance guard of purchasing agents and physics teachers. The lander travels in the same direction with a speed of $0.800c$ relative to the mother ship. As observed on the Earth, the spaceship is 0.200 ly from the Earth when the lander is launched. (a) What speed do the Earth observers measure for the approaching lander? (b) What is the distance to the Earth at the time of lander launch, as observed by the aliens? (c) How long does it take the lander to reach the Earth as observed by the aliens on the mother ship? (d) If the lander has a mass of 4.00×10^5 kg, what is its kinetic energy as observed in the Earth reference frame?
64. A physics professor on the Earth gives an exam to her students, who are in a spacecraft traveling at speed v relative to the Earth. The moment the craft passes the professor, she signals the start of the exam. She wishes her students to have a time interval T_0 (spacecraft time) to complete the exam. Show that she should wait a time interval (Earth time) of

$$T = T_0 \sqrt{\frac{1 - v/c}{1 + v/c}}$$

before sending a light signal telling them to stop. (*Suggestion:* Remember that it takes some time for the second light signal to travel from the professor to the students.)

65. Spacecraft I, containing students taking a physics exam, approaches the Earth with a speed of $0.600c$ (relative to the Earth), while spacecraft II, containing professors proctoring the exam, moves at $0.280c$ (relative to the Earth) directly toward the students. If the professors stop the exam after 50.0 min have passed on their clock, how long does the exam last as measured by (a) the students (b) an observer on the Earth?
66. Energy reaches the upper atmosphere of the Earth from the Sun at the rate of 1.79×10^{17} W. If all of this energy were absorbed by the Earth and not re-emitted, how much would the mass of the Earth increase in 1.00 yr?
67. A supertrain (proper length 100 m) travels at a speed of $0.950c$ as it passes through a tunnel (proper length 50.0 m). As seen by a trackside observer, is the train ever completely within the tunnel? If so, with how much space to spare?
68. Imagine that the entire Sun collapses to a sphere of radius R_g such that the work required to remove a small mass m from the surface would be equal to its rest energy mc^2 . This radius is called the *gravitational radius* for the Sun. Find R_g . (It is believed that the ultimate fate of very massive stars is to collapse beyond their gravitational radii into black holes.)
69. A particle with electric charge q moves along a straight line in a uniform electric field \mathbf{E} with a speed of u . The electric force exerted on the charge is $q\mathbf{E}$. The motion and the electric field are both in the x direction. (a) Show that the acceleration of the particle in the x direction is given by

$$a = \frac{du}{dt} = \frac{qE}{m} \left(1 - \frac{u^2}{c^2}\right)^{3/2}$$

- (b) Discuss the significance of the dependence of the acceleration on the speed. (c) **What If?** If the particle

starts from rest at $x = 0$ at $t = 0$, how would you proceed to find the speed of the particle and its position at time t ?

70. An observer in a coasting spacecraft moves toward a mirror at speed v relative to the reference frame labeled by S in Figure P39.70. The mirror is stationary with respect to S. A light pulse emitted by the spacecraft travels toward the mirror and is reflected back to the craft. The front of the craft is a distance d from the mirror (as measured by observers in S) at the moment the light pulse leaves the craft. What is the total travel time of the pulse as measured by observers in (a) the S frame and (b) the front of the spacecraft?




Figure P39.70

71. The creation and study of new elementary particles is an important part of contemporary physics. Especially interesting is the discovery of a very massive particle. To create a particle of mass M requires an energy Mc^2 . With enough energy, an exotic particle can be created by allowing a fast moving particle of ordinary matter, such as a proton, to collide with a similar target particle. Let us consider a perfectly inelastic collision between two protons: an incident proton with mass m_p , kinetic energy K , and momentum magnitude p joins with an originally stationary target proton to form a single product particle of mass M . You might think that the creation of a new product particle, nine times more massive than in a previous experiment, would require just nine times more energy for the incident proton. Unfortunately not all of the kinetic energy of the incoming proton is available to create the product particle, since conservation of momentum requires that after the collision the system as a whole still must have some kinetic energy. Only a fraction of the energy of the incident particle is thus available to create a new particle. You will determine how the energy available for particle creation depends on the energy of the moving proton. Show that the energy available to create a product particle is given by

$$Mc^2 = 2m_p c^2 \sqrt{1 + \frac{K}{2m_p c^2}}$$

From this result, when the kinetic energy K of the incident proton is large compared to its rest energy $m_p c^2$, we see that M approaches $(2m_p K)^{1/2}/c$. Thus if the energy of the incoming proton is increased by a factor of nine, the mass you can create increases only by a factor of three. This disappointing result is the main reason that most modern accelerators, such as those at CERN (in Europe), at Fermilab (near Chicago), at SLAC (at Stanford), and at DESY (in Germany), use *colliding beams*. Here the total momentum of a pair of interacting particles can be zero. The

center of mass can be at rest after the collision, so in principle all of the initial kinetic energy can be used for particle creation, according to

$$Mc^2 = 2mc^2 + K = 2mc^2 \left(1 + \frac{K}{2mc^2} \right)$$

where K is the total kinetic energy of two identical colliding particles. Here if $K \gg mc^2$, we have M directly proportional to K , as we would desire. These machines are difficult to build and to operate, but they open new vistas in physics.

- 72. A particle of mass m moving along the x axis with a velocity component $+u$ collides head-on and sticks to a particle of mass $m/3$ moving along the x axis with the velocity component $-u$. What is the mass M of the resulting particle?
- 73. A rod of length L_0 moving with a speed v along the horizontal direction makes an angle θ_0 with respect to the x' axis. (a) Show that the length of the rod as measured by a stationary observer is $L = L_0[1 - (v^2/c^2) \cos^2 \theta_0]^{1/2}$. (b) Show that the angle that the rod makes with the x axis is given by $\tan \theta = \gamma \tan \theta_0$. These results show that the rod is both contracted and rotated. (Take the lower end of the rod to be at the origin of the primed coordinate system.)
- 74. Suppose our Sun is about to explode. In an effort to escape, we depart in a spacecraft at $v = 0.800c$ and head toward the star Tau Ceti, 12.0 ly away. When we reach the midpoint of our journey from the Earth, we see our Sun explode and, unfortunately, at the same instant we see Tau Ceti explode as well. (a) In the spacecraft's frame of reference, should we conclude that the two explosions occurred simultaneously? If not, which occurred first? (b) **What If?** In a frame of reference in which the Sun and Tau Ceti are at rest, did they explode simultaneously? If not, which exploded first?
- 75. A ^{57}Fe nucleus at rest emits a 14.0-keV photon. Use conservation of energy and momentum to deduce the kinetic energy of the recoiling nucleus in electron volts. (Use $Mc^2 = 8.60 \times 10^{-9}$ J for the final state of the ^{57}Fe nucleus.)
- 76.  Prepare a graph of the relativistic kinetic energy and the classical kinetic energy, both as a function of speed, for an object with a mass of your choice. At what speed does the classical kinetic energy underestimate the experimental value by 1%? by 5%? by 50%?

Answers to Quick Quizzes

- 39.1 (c). While the observers' measurements differ, both are correct.
- 39.2 (d). The Galilean velocity transformation gives us $u_x = u'_x + v = 110 \text{ mi/h} + 90 \text{ mi/h} = 200 \text{ mi/h}$.
- 39.3 (d). The two events (the pulse leaving the flashlight and the pulse hitting the far wall) take place at different locations for both observers, so neither measures the proper time interval.
- 39.4 (a). The two events are the beginning and the end of the movie, both of which take place at rest with respect to the spacecraft crew. Thus, the crew measures the proper

time interval of 2 h. Any observer in motion with respect to the spacecraft, which includes the observer on Earth, will measure a longer time interval due to time dilation.

- 39.5 (a). If their on-duty time is based on clocks that remain on the Earth, they will have larger paychecks. A shorter time interval will have passed for the astronauts in their frame of reference than for their employer back on the Earth.
- 39.6 (c). Both your body and your sleeping cabin are at rest in your reference frame; thus, they will have their proper length according to you. There will be no change in measured lengths of objects, including yourself, within your spacecraft.
- 39.7 (d). Time dilation and length contraction depend only on the relative speed of one observer relative to another, not on whether the observers are receding or approaching each other.
- 39.8 (c). Because of your motion toward the source of the light, the light beam has a horizontal component of velocity as measured by you. The magnitude of the vector sum of the horizontal and vertical component vectors must be equal to c , so the magnitude of the vertical component must be smaller than c .
- 39.9 (a). In this case, there is only a horizontal component of the velocity of the light, and you must measure a speed of c .
- 39.10 (a) $m_3 > m_2 = m_1$; the rest energy of particle 3 is $2E$, while it is E for particles 1 and 2. (b) $K_3 = K_2 > K_1$; the kinetic energy is the difference between the total energy and the rest energy. The kinetic energy is $4E - 2E = 2E$ for particle 3, $3E - E = 2E$ for particle 2, and $2E - E = E$ for particle 1. (c) $u_2 > u_3 = u_1$; from Equation 39.26, $E = \gamma E_R$. Solving this for the square of the particle speed u , we find $u^2 = c^2(1 - (E_R/E)^2)$. Thus, the particle with the smallest ratio of rest energy to total energy will have the largest speed. Particles 1 and 3 have the same ratio as each other, and the ratio of particle 2 is smaller.



© 2003 by Sidney Harris